

**Elucidation and Inhibition of Sembung Delan  
(*Sphaeranthus indicus* L.) Leaf Extract against  
Balinese Lontar Destructive Fungi**

By

**I.B.G. Darmayasa, Retno Kawuri and I. Wayan Suanda**

ISSN 2319-3077 Online/Electronic

ISSN 0970-4973 Print

Index Copernicus International Value

IC Value of Journal 82.43 Poland, Europe (2016)

Journal Impact Factor: 4.275

Global Impact factor of Journal: 0.876

Scientific Journals Impact Factor: 3.285


InfoBase Impact Factor: 3.66

J. Biol. Chem. Research

Volume 37 (1) 2020 Pages No. 30-35

**Journal of  
Biological and  
Chemical Research**

An International Peer Reviewed / Referred Journal of Life Sciences and Chemistry



**Indexed, Abstracted and Cited in various International and  
National Scientific Databases**

Published by Society for Advancement of Sciences®



**I.B.G. Darmayasa**

[http:// www.sasjournals.com](http://www.sasjournals.com)

[http:// www.jbcr.co.in](http://www.jbcr.co.in)

[jbiolchemres@gmail.com](mailto:jbiolchemres@gmail.com)

RESEARCH PAPER

Received: 26/01/2020

Revised: 28/02/2020

Accepted: 01/03/2020

## **Elucidation and Inhibition of Sembung Delan (*Sphaeranthus indicus* L.) Leaf Extract against Balinese Lontar Destructive Fungi**

**I.B.G. Darmayasa, \*Retno Kawuri and \*\*I. Wayan Suanda**

**Microbiology Laboratory Faculty of Mathematics and Natural Sciences, Udayana  
University, Denpasar, Indonesia**

**\*Microbiology Laboratory Faculty of Mathematics and Natural Sciences, Udayana  
University, Denpasar, Indonesia**

**\*\*Prodi pendi. Biologi FPMIPA IKIP PGRI Bali, Indonesia**

### ABSTRACT

*Lontar is a manuscript written on lontar leaves containing Balinese script that is passed down from generation to generation. The presence of fungal activity can damage the lontar bali in storage. Efforts to control fungus on lontar bali need to be done. One of them is by utilizing natural ingredients from plants. Preliminary research obtained 7 types of fungi isolated from 6 different locations in the province of Bali, namely *Penicillium restrictum*, *Aspergillus fumigatus*, *Mucor racemosus*, *Candida krusei*, *Aspergillus niger*, *Fusarium sp.* and *Rhodoterolamucilaginosus*. This study aims to determine the class of compounds contained in the extract of Sembung Delan leaf (*Sphaeranthus indicus* L.) and its inhibition on 7 types of fungi that have been isolated from lontar Bali. In vitro testing of the inhibition of the Sembung Delan leaf extract on each fungal isolated on lontar bali was carried out by the diffusion well method. The compound group contained in the leaf extract of Sembung Delan was known through phytochemical tests. The results showed that all types of fungi isolated in Balinese palm oil were able to be inhibited by crude extracts of Sembung Delan leaves with different barriers. Phytochemical tests are known that the sembung Delan leaves contain alkaloid, terpenoids, phenolic, saponin, flavonoid and tannin compounds.*

*Keywords: Lontar bali, Fungi, Sphaeranthus indicus and Extract.*

### INTRODUCTION

Lontar is one cultural heritage in form of manuscripts containing Balinese scripts which have been handed down from generation to generation. The manuscripts written on palm leaves are considered classic or ancient which tends to be sacred and religious (Geriani, 2010, Sedana et al., 2013). The environment conditions with high humidity might cause lontar to be easily damaged. In addition to insects, the existence of microorganisms such as fungi became another cause of the damage. Rontar leaves from rontar plants (*Borassus flabellifer*) as raw material for making lontar are mostly composed of cellulose components which are good substrates for fungal growth. According to Yosmar et al. (2015), cellulose is a carbohydrates making up the cell walls of plants that are easily damaged by cellulase enzyme.

Sancana (2014) stated that lontar manuscript as a cultural object does not have a strong resistance to environmental influences. Furthermore, there were reportedly several groups of fungi as a cause of damage to the Balinese lontar, namely: *Aspergillus*, *Penicillium* and *Fusarium*. In the initial research, there were 7 types of fungi found to be isolated from Balinese lontar in different locations, namely: 2 locations in Denpasar City (GriyaBalun and GriyaBatukandik); 2 locations in Tabanan Regency (Griya Kediri and GriyaSunantaya) and 2 locations in Gianyar Regency (GriyaUbud 1 and GriyaUbud 2). The types of fungi found were *Penicilliumrestricum*, *Aspergillus fumigatus*, *Mucorramosus*, *Candida krusei*, *Aspergillus niger*, *Fusarium* sp. and *Rhodoterolamucilaginoso*. The growth of fungi in Balinese lontar needs to be managed so that the damage can be prevented. In this study, using natural materials as a source of raw materials in designing the formulations to inhibit the growth of fungi that could damage Balinese lontar.

SembungDelan leaves (*Sphaerantusindicus* L.) are plants which grow agricultural land that have a potential to control the growth of fungi. A research from Darmayasa (2002) proved that the extract of this leaf was able to inhibit *Alternariasp.*, *Gleosporiumsp* and *Phytophthorasp.* fungi in *in-vitro*. Furthermore, Darmayasa (2014) reported that the extract of SembungDelan was able to inhibit the growth of *Phytophthorainfestanfungi* which is the cause of leaf blight in potato plants. Based on those researches, a deeper research on the ability of SembungDelan leaves to inhibit the growth of fungi in lontar Bali needs to be conducted.

## MATERIAL AND METHODS

### Time and Place of the Research

This research is conducted from May to September, 2019 in the Microbiology Laboratory, Faculty of Mathematics and Natural Sciences, Udayana University and a joint Laboratory of the Mathematics and Natural Sciences of Udayana University.

### Method of Extraction

The SembungDelan leaves are washed clean and air dried. Then, blended until become flour consistency. Next, measured to 100grammacerated with 1 liter of methanol at room temperature for 72 hours. The maceration results were then filtered. The filtrate obtained was evaporated with a vacuum rotary evaporator at 40°C to separate the solvent. The crude extract obtained was used for further analysis (Harborne, 1996).

### The Inhibitory Test of SembungDelan (*S. indicus* L.) Crude Extract on Balinese Lontar Destructive Fungi

The inhibitory test of SembungDelan (*S. indicus*L.) on Balinese lontar destructive fungi used the diffusion well method (Hewitt and Vincent, 1989). First, prepare each fungal suspension that had been isolated from the Balinese lontar by media *Potato Dextrosa Agar*, poured as much as 15 mL into a sterile petri dish. Next, implant the fungal colonies taken from culture stock of the sloping media right in the middle of the petri dish. After incubation time of 4 days at room temperature, the growing fungi were then rinsed using 5 mL of sterile water in order to obtain a suspension of mold spores. The density of mold spores was calculated using *NouemHaemocytometer*.

Each suspension of mold spores was taken as much as 200  $\mu$ L. Next, put into a sterile Petri dish then poured PDA media and homogenized in order to obtain an even growth of fungi. After the mixture of PDA media and isolate was frozen up, several wells with a diameter of 0,5mm were made on that media. As much as 40  $\mu$ L crude extract of SembungDelan leaves was put into the well. However, for the negative control on the other wells, only methanol was being deposit. All of those treatments were repeated 3 times. The inhibitory of the crude extract of SembungDelan leaves on the tested fungi was determined by measuring the diameter of the clear zone around the well. The measurements were conducted using calipers for 4 times and then being averaged. The crude extract of SembungDelan leaves which has shown a potential as a vegetative fungicide is given a concentration of 5% (b/v) by dilution using the following formula:

$$V_1 \cdot M_1 = V_2 \cdot M_2$$

Keterangan:

Note:

$V_1$ : initial volume  $V_2$ : expected volume  $M_1$ : initial concentration  $M_2$ : expected concentration

### **The Phytochemical Test of the crude extract of Sembungdelan leaves (*Sphaeranthusindicus*L.)**

The Phytochemical test of the crude extract of SembungDelan leaves being conducted were the test for alkaloids, flavonoids, phenolics, steroids, terpenoids, and saponins.

#### **Test for Alkaloids**

The sample of Sembungdelan leaf extract was dissolved into 10 ml of ammonia chloroform, then added 0.5 mL of H<sub>2</sub>SO<sub>4</sub> and homogenized, after that placed until two layers were formed which were pulp in the lower layer and clear extract in the upper layer. The top layer was taken, then dripped with meyer reagent as much as 1 drop. If the extract turn out to be positive to contain alkaloids, it would form intosediment. (Harborne, 1996).

#### **Test for Flavanoids and Phelonics**

Test for flavonoids and phelonics in Sembungdelan extract was conducted by dissolving the extract in to 70% of ethanol and being heated up. Then, the extract was filtered to obtain the filtrate, and the result of the filtrate would be placed onto a drip plate and added with Mg and HCL to test for flavonoids, and added with FeCL<sub>3</sub> to test for the phenolics. The change of color on the extract would be red to show that it contained flavonoids, while the formation of a green to purplish green ring on the extract would signify the phenolic extract (Harborne, 1996).

#### **Test for Steroids and Terpenoids**

Test for steroids and terpenoids in Sembungdelan extract was conducted by adding a sample of Sembungdelan leaf extract into chloroform and heated up for 10 minutes. After being heated up, the sample was placed on a drip plate and then added with Lb reagent (concentrated antridic acetic acid + concentrated H<sub>2</sub>SO<sub>4</sub>). If the extract turned out to be positive to containterpenoids, the color would changeinto red, pink or violet in the SembungDelan extract, and if the sample turned out to be positive to contain steroids, the color would changeinto green or purple in the sample (Harborne, 1996).

#### **Test for Saponins**

Test for saponins in the sembungdelan extract was conducted by dissolving as much as 0,1 g Sembungdelan leaf extract into 5 mL of hot aquadest and shaken for 10 seconds. If the extract turned into a stable froth or foam for 10 minutes, the extract must be positive to contain saponins (Harborne, 1996).

### **The Analysis of the Data**


The obtained data in this research were in qualitative, which is showed in form of pictures or graphics, and in quantitative, which is analyzed using the Analysis of Variance (ANOVA). If the obtained data showed a significant difference at the test level of 5% (P <0.05), it would be proceeded into Duncan test.

## **RESULT AND DISCUSSION**

The crude extract of SembungDelan leaves at the concentration of 0,01g/20 μL in *in vitro* was able to inhibit the growth of fungi in Balinese lontar. This matter is shown by the presence of the diameter of the inhibitory zone that formed around the diffusion well. The diameter of the inhibitory zone of each fungus tested was different. The biggest resistance diameter occurred in the testing of SembungDelan crude extracts on the fungus *Aspergillus niger* was 21 mm. The smallest resistance obtained in the test of the fungus *Rhodoterulamucilaginoso* was with a diameter of the inhibitory zone of 10 mm, more explanation is presented in Table 2.

All types of fungi that have been isolated from Balinese lontar in different places in *in viro* were able to be inhibited by the crude extract of SembungDelan leaves at the concentration of 0,01g/20 μL. In this study, the inhibitory had by the crude extract of SembungDelan leaves were varied. The difference of the inhibitory was thought to be closely related to the structure and characteristics of the isolated fungi. The biggest inhibition based on the measurement of the diameter of the inhibitory zone occurred in the *Aspergillus niger* fungus, which was 21 mm. The smallest inhibition was found in the testing of *Rhodoterulamucilaginoso* fungus with a diameter of 10 mm in the inhibition zone. According to Nester *et al.* (2007), there appears to be a difference of inhibitory from one substance tested in *in vitro* is determined by the type, concentration of the test compound, and type and concentration of the microbes tested.

A similar research conducted by Darmayasa (2014) reported that Sembung Delan leaves were able to inhibit the growth of *Phytophthora infestans* fungi which is the cause of leaf blight in potato plants. Similarly to the research conducted by Mhetreet *al.*(2006) which showed that Sembung Delan plants were able to inhibit the growth of *Stappyllococcus aureus*, *Eschericia coli*, *Fusarium sp.*, and *Penicillium pinophilum*. The formation of the clear zone diameter around the diffusion well in this study proved that there was an active substance that functioned as an antifungal contained in the crude extract of Sembung Delan leaves.



Note:

- 1 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Candida krusei*
- 2 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Aspergillus fumigatus*
- 3 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Mucor racemosus*
- 4 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Aspergillus niger*
- 5 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Rhodotterula mucilaginosa*
- 6 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Fusarium sp*
- 7 sign (a) control (b) Diameter of inhibitory zone of Sembungdelan leaf extract against *Penicillium restrictum*

The phytochemical test of Sembungdelan leaf extract, it turned out to be positive to contain alkaloids, terpenoids, phenolics, saponins, flavonoids and tannins. However, the test for steroids showed a negative result. Qualitatively, the most abundant compound in SembungDelan leaf extract was terpenoids, proven from the test results giving a prominent reddish color. The least amount of compounds in SembungDelan leaf extract was alkaloids and tannins. More details are presented in Table 3.

**Table 2. The diameter of the inhibitory zone of SembungDelan leaves (*S. indicus*L) crude extract on the fungi in Balinese lontar.**

No	Type of Fungus	Inhibitory of SembungDelan( <i>Sphaerantus indicus</i> L.)crude extract (Ø mm)
1	<i>Penicillium restrictum</i>	18±0,24
2	<i>Aspergillus fumigatus</i>	18±0,25
3	<i>Mucor racemosus</i>	20±0,15
4	<i>Candida krusei</i>	19±0,24
5	<i>Aspergillus niger</i>	21±0,15
6	<i>Fusarium</i> sp	19±0,24
7	<i>Rhodoterulamucilaginoso</i>	10±0,00

**Table 3. Compound groups of SembungDelan leaf extract based on the Phytochemical test.**

No	Compound Group	result	Unit
1	Alkaloids	+	white sediment
2	Steroids	-	greenish color
3	Terpenoids	+++	reddish color
4	Phenolics	++	greenish color
5	Saponins	++	foam
6	Flavonoids	++	reddish color
7	Tannins	+	bluish color

Note:

+++ :give sediment/a lot of color/prominent

++ : give sediment/medium color

+ :give sediment/enough color

- : no sediment/color

Based on the phytochemical test, SembungDelan plants contained several compound groups namely alkaloids, terpenoids, phenolics, saponins, flavonoids and tannins. The presence of the compound groups has been reported by Tiwari and Khosa (2009) in SembungDelan plants in the mainland of India, which contained *flavonoid glikosida, flavonoid C-glikosida, isoflavon, sterol, alkaloid, asam amino, 2-hydroxycostic acid, asamilicic*, essential oil which has methyl chavicol, d-cadinene,  $\alpha$ -ionone, p-methoxycinnamaldehyde, eugenol, taninns and steroids. In this study, test for steroids has turned out to be negative. The difference in geographical condition was assumed to be the cause of the absence of steroid compounds in SembungDelan plants in mainland of Indonesia. The essential oil that is contained in SembungDelan leaves has the characteristic of being an antifungal against *Trichoderma viride, Rhizopus nodosus, Aspergillus niger, Trichophyton rubrum* and *Curvularia parasadii* (Mane and Badole, 2013). Emaniet *et al.* (2017) successfully isolated the compound *5 $\alpha$ -hydroperoxy-7 $\alpha$ -hydroxy-isosphaerantholide* and *(11 $\alpha$ ,13-dihydro-7 $\alpha$ -hydroxyfrullanolide-13-yl)-adenine (2)* from its flowers. However, it is still cannot be proven whether the isolated compound could be used as antifungal. It is in contrast with Bhuwanet *et al.* (2016) which stated that the extract of SembungDelan plants (*S. indicus*L.) had a very broad biological activity. Selvanayagamet *et al.* (2005) said that *S. indicus*L. were indigenous plants which grew on agricultural land, and is widely used as a medicine for dysentery, diseases of the uterus and vagina, urinary tract infections as well as a control of worms and pests. More over, stated in their research that *S. indicus*L. contained crystal with a structure of 3 $\alpha$ -Hydroxy-3,5 $\alpha$ ,9-trimethyl 1,3,3 $\alpha$ ,4,5,5 $\alpha$ ,6,7,8, 9b, decahydro-2naphtha [1,2c]-imidazole-2-one Monohydrat. Reported by Sangeetha *et al.* (2010) that the extract of *S. indicus* L. using ethanol solvent, chloroform and petroleum were able to inhibit microbes that are the causes of several infection diseases in humans, such as *Candida albicans, Cryptococcus neoformans, Salmonella typhy, Klebsiellasp., Pseudomonas aeruginosa*, and *E. coli*.

## CONCLUSION

The Extract of SembungDelan leaves (*S. indicus*L.) with methanol solvent was able to inhibit the growth of *Penicilliumrestricum*, *Aspergillus fumigatus*, *Mucorracemosus*, *Candida krusei*, *Aspergillus niger*, *Fusarium*sp. and *Rhodotorulamucilaginosafungi* which were isolated from Balinese lontar. The compound groups contained in the SembungDelan leaf extract in this study were alkaloids, terpenoids, phenolics, saponins, flavonoids and tannins.

## ACKNOWLEDGMENTS

I would like to thank the Head of LPPM UniversitasUdayana for providing the research fundings through PUU 2019 with SK. NOMOR 357 / UN14 / HK /2019 and also for the people who have been supporting the process of writing this paper.

## REFERENCES

- Al-Enazi, N.M., A.S. Awaad, M.R. Al-Othman, N.K. Al-Anazi, and S.I. Alqasoumi (2017). Isolation, identification and anti-candidal activity of filamentous fungus from Saudi Arabia soil. *Saudi Pharmaceutical Journal*. x(xx)
- Deligios, M, C. Fraumene, M. Abbondio, I. Mannazzu, A. Tanca, M.F. Addis and S. Uzzau (2005). Draft Genome Sequence of *Rhodotorula mucilaginoso*, an Emergent Opportunistic Pathogen. *Journal ASM org*. 3(2): 1-15.
- Gandjar, I., R.A. Samson, K.V.T. Vermiolen, A. Oetari, and I. Santoso (1999). *Pengenalan Kapang Tropik Umum*. Universitas Indonesia. Depok.
- Geriani, A. (2010). Lontar: Tradisi Hidup dan Lestari di Bali. *Media Perpustakaan*. 17(1): 39-45.
- Hai, H.D. (2016). Toddy Palm (*Borassus flabellifer*). Diakses di worldwide fruits padatanggal 22 April 2018.
- Hachem, R., H. Hanna, D. Kantoyiannis, Y. Jiang and I. Raad (2018). *Candida glabrata* and *Candida krusei* as the Leading Causes of Candidemia in Hematologic Malignancy. American Cancer Society. Published Online 15 April 2008 in Wiley Inter Science. ([www.interscience.wiley.com](http://www.interscience.wiley.com)).
- Palmweb (2017). Palmweb: Palms of the World Online. Diakses di <http://www.palmweb.org>. padatanggal 12 Juli 2018.
- Pitt, J. I. and A. D. Hocking (1997). *Fungi and Food Spoilage*. Second Edition. Printed in Great Britain at the University Press, Cambridge.
- Sancana, A. (2014). Kajian Efektivitas Teknik dan Bahan Konservasi pada Lontar di Bali. *Jurnal Konservasi Cagar Budaya Borobudur*. 8(2):11-29.
- Sedana, I.N., N.A. Damayanti, dan U.L.S. Khadijah (2013). Preservasi Berbasis Kearifan Lokal (Studi Kasus Mengenai Preservasi Preventif dan Kuratif Manuskrip Lontar sebagai Warisan Budaya di Kabupaten Klungkung Bali). *Jurnal Kajian Informasi & Perpustakaan* 1(1):91-105.
- Setiani, K. (2018). Potensi Ekstrak Filtrat *Trichoderma* sp. Dalam Menghambat Pertumbuhan *Aspergillus flavus* pada Lontar Bali di Griya Gede Siwa Manggis Manuba Desa Nyanglan Kecamatan Banjarangkan Kabupaten Klungkung, Bali (*Undergraduate thesis*). Jur. Biologi FMIPA Unud.
- Sirait, B., N.S. Napitupulu, A.Y. S. Simanjorang and A.I. Manurung (2018). Preliminary Research the Effect of NAA and BAP on Callus of Benana (*Musa acuminata* L.) With Addition of Fusarium Wilt Disease from *Fusarium oxysporum* sp. cubense (Foc). *Jurnal Darma Agung* 26(1):628-632.
- Sukamaluddin, Mulyadi, Dirawan, G. D., Amir, F. and Pertiwi, N. (2016). Conservation Status of Lontar Palm Trees (*Borassus flabellifer* Linn) In Jeneponto District, South Sulawesi, Indonesia. *Journal of Tropical Crop Science*, 3 (1): 28-33.
- Talantan, V.M., Marina, O. Lambui and I.N. Suwastika (2018). Uji Aktivitas Selulase Dari Jamur Selulolitik Asal Tanah Danau Kalimpa'a Sulawesi Tengah.
- Yosmar, R., N. Suharti, dan R. Rasyid (2013). Isolasi dan Uji Kualitatif Hidrolisis Jamur Penghasil Enzim Selulase dari Tanah Tumpukan Ampas Tebu. *Jurnal Farmasi Andalas*. 1(1):5-12.

---

Corresponding author: I.B.G. Darmayasa, Microbiology Laboratory Faculty of Mathematics and Natural Sciences, Udayana University, Denpasar, Indonesia  
Email: [darm\\_aponk@yahoo.co.id](mailto:darm_aponk@yahoo.co.id)