# Pesticide Residues in Vegetable and Fruit Samples from Andhra Pradesh, India By

A. Harinathareddy, N.B.L. Prasad and K. Lakshmi Devi

ISSN 0970-4973 (Print) ISSN 2319-3077 (Online/Electronic)

Index Copernicus International Value IC Value of Journal 4.21 (Poland, Europe) (2012) Global Impact factor of Journal: 0.587 (2012) Scientific Journals Impact Factor: 2.597

J. Biol. Chem. Research Volume 31 (2) 2014 Pages No. 1005-1015

# Journal of Biological and Chemical Research

(An International Journal of Life Sciences and Chemistry)

Indexed, Abstracted and Cited in Various National and International Scientific Databases of the World

Published by Society for Advancement of Sciences®



Received: 12/06/2014 Revised: 25/08/2014

RESEARCH PAPER Accepted: 01/09/2014

# Pesticide Residues in Vegetable and Fruit Samples from Andhra Pradesh, India A. Harinathareddy, \*N.B.L. Prasad and \*\*K. Lakshmi Devi

Department of Bio technology, JNTUA, Anantapuramu, Andhra Pradesh, India OTRI, JNTUA, Anantapuramu, Andhra Pradesh, India

\*\*Department of Biochemistry, S.K. University, Anantapuramu, Andhra Pradesh, India

# ABSTRACT

Vegetable samples of brinjal, okra, green chilli, cabbage, Cauliflower, Tomato and Capsicum from the farmers' field Fruit samples like Banana, Pomegranate, Sweet Orange, Guava, and Grapes collected from Market in five regions of Andhra Pradesh. Were tested for the presence of organochlorine (OC), organophosphorus (OP), and synthetic parathyroid (SP) compounds using a gas chromatograph equipped with electron capture and Thermo sensitive detectors of the samples tested, 36.60 % were found to have pesticide residues. Among the OC compounds,  $\alpha$ -endosulfan,  $\beta$ -endosulfan, were detected in 3.33 % of the samples with residues. These were taken from Cabbage and Brinjal samples. SP compound residues, such deltamethrin were detected in 1.66 % of the samples with residues. OP compound residues such as chlorpyrifos, profenofos, ethion, dimethoate, Malathion, Quinalphos and methyl parathion were found in 22 % of the samples with residues, which were taken from all vegetable and fruit samples of the positive samples, 15.3 % were found to contain residues exceeding the prescribed maximum residue limit. The average pesticide residue content across all the vegetable samples was 0.03 ppm, with values ranging from 0.03 to 1.18 ppm. Multiple residues of more than one compound were detected in 13.33 % of samples containing residues.

*Keywords: Pesticide residues, Vegetables and Fruits, Organochlorine (OC), Organophosphorus (OP), Synthetic Parathyroid (SP) and Gas Chromatography.* 

Published by Society for Advancement of Science®

## INTRODUCTION

Vegetables are an important part of balanced diet providing many essential minerals and vitamins. Besides, higher intakes of vegetables are associated with healthier lives including lower risks of cancer, cardiovascular diseases and obesity. But the yield and quality of vegetables are affected by pests and diseases infestation during production and storage necessitating the use of pesticides. The pesticide use has increased over the years for effective controlling of a pest complex of vegetable crops due to rapid action and less manpower requirement. However, injudicious use of these compounds often leads to the presence of pesticide residues in harvested produce in amounts exceeding prescribed maximum residue limits (MRL), which may pose serious health risks as some of the vegetables are eaten raw or undercooked.

India is one of the largest agricultural pesticide consumers worldwide and is the second largest manufacturer of pesticides in Asia (Mathur 1999). There is a sequential rise in the production and consumption of pesticides in India during last three decades. The consumption pattern of pesticides differs from rest of the world, as in India, 76 % of total pesticide consumption is insecticides as against 44 % worldwide. Of the total pesticide usage 45 % goes to cotton crop followed by paddy and wheat. While only 10-12 % of total pesticides are used for fruits and vegetables (Kumari et al. 2003; Bhattacharyya et al. 2009). In Andhra Pradesh, vegetable cultivation has increased in recent years because of greater market demand, higher returns, and an increased awareness among farmers. This has led to large stretches of vegetablegrowing areas with intensive management practices. Such intensive cultivation, coupled with the evailing humid tropical climate, has resulted in increased pest and disease incidence. Nearly 75 to 80 % of total pesticide consumption of 5175.4 MT is used for vegetable crops alone (Andaman and icobar Administration 2009) as against the national average of 10-12 %. As the vegetables are grown in 449.2 ha only, the per capita pesticide consumption in the vegetablegrowing areas of Andhra Pradesh (3,586 g) far exceeds the national average of 450 g (Bhattacharyya et al. 2009).

There are several reports on environmental contamination and long-term health implications resulting from the intensive use of pesticides in agriculture, animal husbandry, and public health programs (lgbedioh 1991; Waliszevski et al. 1996; Bhanti and Taneja 2007; Chen et al. 2011). As fruits and vegetables are the most frequently consumed food group, which makes up 30 % of Food consumption on mass basis (WHO 2003), the presence of pesticides in vegetables is a moreserious health concern, since they are consumed raw, without much processing and storage (Kalra and Chawla 1985; Kumari et al. 2003). Because of which, they are expected to contain higher levels of pesticide residues than other food groups of plant origin. Aside from causing acute and chronic toxicity, they can also affect the immune system and cause arthritis, carcinogenicity, reduced fertility, increased cholesterol, high infant mortality, etc. (Singh 2001). Thus, the monitoring of pesticide residues is essential to regulate the application of pesticides to food crops especially vegetables so as to ensure healthy products to the consumers.

Several studies have been reported on the contamination of vegetables in India (Agnihotri 1999; Mukherjee 2003; Kumari et al. 2004; Bhanti and Taneja 2005; Mandal and Singh 2010; Chowdhuryetal. 2011) and world (Adeyeye and Osibanjo 1999; Amoah et al. 2006; Nishina et al. 2010; Bempahetal. 2012). The intensive cultivation of vegetables has gained momentum in recent years with excessive pesticide usage due to increased market demand. As the presence of pesticide residues in food produce is a serious concern and no data are available on the levels of pesticide residues in vegetables, a Study was conducted to ascertain the presence of pesticide residues in major vegetables.

# MATERIAL AND METHODS

Andhra Pradesh's GDP for 2011 was approximately ₹ 5,67,636 lakh crore, placing it third among the states. The state ranks second in terms of overall Gross State Product among all the states of the Indian Union. In terms of per capita GSDP the state compares very favorably with other large states.

In the Agriculture has been the chief source of income for the state's economy. Andhra Pradesh is an exporter of many agricultural products. Four important rivers of India, the Godavari, Krishna, Penna, and Thungabhadra flow through the state, providing irrigation. Rice, sugarcane, cotton, Chili pepper, mango, and tobacco are the local crops. Recently, crops used for vegetable oil production such as sunflower and peanuts have gained favor. There are many multi-state irrigation projects in development, including Godavari River Basin Irrigation Projects and Nagarjuna Sagar Dam. In Andhra Pradesh the total area under vegetables is 2.36 lakh hectares with an annual production of 3.31 million tones. The per capita consumption is guite low. The present production is not adequate to meet the present demand of 7.70 million tones. To provide nutritional security and to fill up the gaps it is guite essential to increase the production and productivity levels of vegetables. During 2004-05, area of 23,000 hectares (Kharif 9199 hac and Rabi & Summer 13801 hac) is proposed to be covered under hybrid vegetable seeds with an outlay of Rs.376.75 lakhs. The major vegetable-growing areas in these Andhra pradesh are Kuronool, Chittor, Prakasam, Ranga Reddy, Krishna, Warangal, nalgonda. Brinjal, okra, cauliflowers, cabbages, tomatoes, chilies, and cucurbits are the major vegetables grown. Of the total pesticides used, approximately 75–80 % is applied for vegetable cultivation alone. The vegetable cultivation is practiced in an open-field condition under conventional management practices. The vegetable growers are only small to medium farmers with average land holding not exceeding 2 ha. The major pesticides used on the islands are cypermethrin, malathion, dimethoate, quinalphos, chlorpyrifos, endosulfan, phorate, and carbofuran (Jeya Kumar et al. 2008)

Fruit crops occupy major portion of area coverage i.e. 7.60 lakh hectares in the state. The major crops are mango, citrus, banana, cashew, cocoa and plantation crops. During the year 2004-2005 increase in production, area and productivity is to be achieved through technology and management practices. Capacity building through training is also a thrust area.

Geographically, Andhra Pradesh is composed of most of the eastern half of the Deccan plateau and the plains to the east of the Eastern Ghats. Andhra Pradesh is divided into three regions. The northern part of the plateau is the Telangana region and the southern part is known as Rayalaseema. These two regions are separated by the River Krishna. The third region is Coastal Andhra. The plains to the east of Eastern Ghats form the Eastern coastal plains. The Eastern Ghats are discontinuous and individual sections have local names. The Kadapa Basin, formed by two arching branches of the Eastern Ghats is a mineral-rich area. The coastal plains are for the most part delta regions formed by the Godavari, Krishna, and Penner rivers. The Eastern Ghats are a major dividing line in the state's geography. The Ghats become more pronounced towards the south and extreme north of the coast. The Eastern Ghats give way to the Deccan Plateau, where shrub vegetation is more common. Most of the coastal plains are put to intense agricultural use. The west and southwest parts of Andhra Pradesh have semi-arid conditions.

The multiresidue extraction was done following a modified QuEChERS method (Anastassiades et al. 2003), where the samples are cut into pieces and macerated with a mixer grinder. A 15-g macerated sample is then weighed into a 50-mL centrifuge tube along with 30 mL acetonitrile and shaken well for 1 min in a vortex shaker. The supernatant is transferred quantitatively to a second centrifuge tube containing 4 g anhydrous MgSO4 and 1 g NaCl. The tube was then shaken vigorously again for 1 min in a vortex shaker and centrifuged at 3,000 rpm for 5 min. A8-mL extract of this sample is added to a 15-mL centrifugetube that contains 1.2 g anhydrous MgSO4 and0.4g primary secondary amine, shaken vigorously for 1 min in the vortex shaker, and centrifuged at 3,000 rpm for 5 min. A 2-mL extract sample is pipette out into a glass tube and the solvent is evaporated usinga Turbo Vap evaporator at 30 °C using 10 psi N2 gas over15 min. The residues are reconstituted using n-hexanefor final GC and GC/MS analysis.

The detection and quantification of different pesticide compounds was carried out by injecting 1  $\mu$ L of the extract into the gas chromatograph (GC-Shimadzu GC-2010) equipped with both an electron capture detector and a flame thermionic detector. The residue levels were estimated by comparing the peak areas of the samples to those of standards run under identical conditions. Any detected residues were confirmed using GC/MS (GC/MS-TQD Brukers) to avoid the misinterpretation of any results. Three vegetable samples with no residues were spiked with 0.01 ppm levels of organochlorines, 0.05ppm for organophosphorus, and 0.1 ppm for Synthaticpyrethroids compounds and separately analyzed following the same procedure. More than 80 % recovery was obtained for the spiked samples and results were reported as such with no correction for recovery rate. The average pesticide residue contents in the vegetable samples were calculated using the equation given by Poulsen et al. (2007).

# **RESULT AND DISCUSSIONS**

Pesticide residues were found in 38.3 % of the total samples analyzed which is lower than the reported contamination levels elsewhere in India (Kumari et al. 2004; Charan et al. 2010) and world (Hjorth et al. 2011; Bempah et al. 2012). In 33.3 % of the samples, residues were found at or below MRL, while in 5.0 % samples, residues are above the MRL (PFA 1954). Among individual vegetables, green chillies, capsicum, Tomato, Okra, Brinjal, and cruciferous vegetables like cabbage and cauliflower showed presence of residues in more than 33.3 % of samples analyzed. In fruits, Grapes, Pomegranate samples showed presence of residues in more than 3.3 % of samples. This trend is in consonance with the reports of higher levels of contamination of Okra (Ashutosh 2010), Tomato (Crentsil Kofi Bempah 2011) and Green chilli (Z Praveen 2005) samples. And also from elsewhere in India (Kumari et al. 2003; Charan et al. 2010). In greenchillies, 16.6 % samples showed residues more than specified MRL, followed by Tomato (33.3 %). In Grape 3.3% samples showed the residues followed by Pomegranate 1.66% samples (Beena Kumati 2006, Mutwakil M.A 2005-09).(Table 1). A total compounds, covering organochlorine (OC), organophosphorus (OP), and synthetic pyrethroid (SP) compounds, were detected in this study. The residues of OP (22 %) and SP (1.66 %) compounds were most frequently found, followed by OC (6.6 %) compounds. The residues of OP compounds were predominant in all vegetable groups indicating their widespread usage. The residues of OC (6.6 %) and OP (45.9 %) compounds were detected most in green chillies followed by Tomato and okra vegetables and in fruits (4.9%) grapes followed by pomegranate. (Table 2). Among the compounds detected, Dimetoate, Profenophos, Quinalphosand Malathion were most frequently found in 41.97 % of positive samples. Dimethoate is the single largest pesticide found with 13.3% of the samples surveyed reporting its use followed by Malathion. Malathion is found 11.66 % of samples. Profenophos and Quinalphosis found 6.66 % of samples. As Malathion has relatively apid degradation with less persistence, generally no residues were found as a residue in food. However, in recent times, predominance of Malathion residues in fruits and vegetables was also reported by several workers (Singh and Kiran Singh 2006; Chen et al. 2011; Lozowicka et al. 2012) in their respective studyarea indicating their excessive use. Though endosulfan was banned for its agricultural use in the Andhra Pradesh since May 2011, the presence of its residues in recent samples indicated its illegal use which might be from previous stock. Its continuous monitoring helps in ensuring tight control measures not only on farmers, but also on pesticide dealers. The concentration of pesticide residue ranged from 0.03 to 1.18 ppm with a mean concentration of 0.192 ppm. The mean values of OP (0.184 ppm) and OC (0.071 ppm) residues in positive samples exceeded the overall mean of 0.192 ppm (Table 3). The mean content of SP compounds was found to be low, which might be due to their less persistence. Samples were tested for the presence of OC compounds, viz.,  $\alpha$ hexachlorocyclohexane (HCH), β- HCH, lindane, delta HCH dicofol, di-chlorodiphenyl trichloroethane (DDT) and metabolites thereof, such as o, p'-DDE,o,p'-DDD,p,p'-DDT, p,p'-DDD, p,p-dichlorodiphenyldichloroethylene (DDE), o,p'-DDE,  $\alpha$ -endosulfan,  $\beta$ -endosulfan, and endosulfansulfate.

Residues of endosulfan T ( $\alpha$ -endosulfan,  $\beta$ -endosulfan, and endosulfan sulfate) weredetected in 3.33 % across all samples having residues. At the time of sampling, it was also observed that the crops were heavily infested with diamond back moth and shoot and fruit borers. Because of its relatively low cost and broad spectrum application, it might be widely used by many producers for pest control including thrips and aphids. Its residue level in contaminated samples ranged from 0.03to 1.18 ppm with average content of 0.192 ppm. The highest mean concentration was detected in Tomato (0.358 ppm) followed by Okra (0.331 ppm), Green chilli (0.112 ppm), and Cabbage (0.162 ppm) (Fig. 4). In majority of the samples (60%), the concentration was found to be less than 33.3 % of specified MRL. The highest level of 0.88 ppm that was found occurred in only one sample of Tomato which exceeded the MRL of 0.2 ppm. Synthetic pyrethroids, such as  $\alpha$ -cypermethrin,  $\lambda$ -cyhalothrin, bifenthrin, fenpropathrin,  $\beta$ -cyfluthrin, deltamethrin, fenvaleratel, fenvalerate II, fluvalinatel, and fluvalinate II, were also investigated in the vegetable samples. Residues deltamethrin weredetected in the positive samples. These vegetable samples were also tested for 15 organophosphorus compounds, including dichlorvos, phorate, dimethoate, methyl parathion, fenitrothion, malathion, chlorpyrifosmethyl, chlorpyrifos, guinalphos, profenofos, ethion, and Phosalone. Of these residues, profenofos, chlorpyrifos, Dimethoate, malathion, Quinalphos, Methyl parathion, Ethion and Phorate were found in 36.6 % of the positive samples. These residues were found in all the vegetable groups with values ranging from 0.03 to 1.18 ppm. Chlorpyrifos was dominant OP compound in Green chilli have 0.35ppm have the above MRL (0.2ppm) followed by Phorate 0.08pmm is above MRL. The concentration of individual compounds varied with mean values ranging from 0.03 to 1.18 ppm (Sajjad Ahmed Baig 2009). The residues were found in fruits ranging from 0.05 to 0.10 ppm. The mean concentration of Dimethoate, Chlorpyrifos, Profenophos, Quinalphos, and Malathion was higher than the average value of all OP compounds Table 5). The residues of more than one compound were found in 13.33 % across the samples and 38.3 % of samples containing residues. Among different pesticides, Dimethoate was the most frequently found along with Malathion, profenofos, Quinalphos, and endosulfan. Besides, the Methyl parathion and Ethion association was found in few samples of Green chilli. Multiple residues were highest in green chilli (8.33 %), followed by Okra. In this study, a distinct pattern of pesticide use is discernable from the maximum number of samples containing different pesticide residues. Accordingly, Profenophos, Chlorpyrifos and Dimethoate compounds were mainly used for green chilli and okra; Quinalphosfor brinjal, Endosulfan and Deltamethrin were used for all vegetables.

#### **Figure Legends**

- 1. Fig-1. In total samples percentage of samples vegetables detected.
- 2. Fig-2. Percent contribution of different vegetables to samples with detected samples.
- 3. Fig-3. In total samples percentage of samples fruits detected.
- 4. Fig-4. Percent contribution of different Fruits to samples with detected samples.

#### Tables

- 1. Table 1: Summary of pesticide residue analysis in vegetables and fruits
- 2. Table 2: Distribution of various pesticide groups among contaminated vegetable groups
- 3. Table 3: Pesticide residue levels in vegetables and fruit samples
- 4. Table 4: Concentration of OCs, OPs and SPs in Vegetable samples



Fig 1. In total samples percentage of samples vegetables detected.



#### Fig 2.Percent contribution of different vegetables to samples with detected samples.



Fig 3. In total samples percentage of samples fruits detected.



Fig. 4. Percent contribution of different Fruits to samples with detected samples.

Table 1.Summar	y of pe	sticide residue	analysis in	vegetable	es and fruits.
----------------	---------	-----------------	-------------	-----------	----------------

Commodity	No of samples analyzed	Samples With detected residues	% total samples	Samples With out detected residues	% total samples	Samples With detected residues Below MRL or without MRL	% total samples	Samples With detected residues above MRL	% total samples
Vegetables	30	20	66.6	10	33.3	17	56.6	3	10
fruits	30	3	10	27	90	3	10	0	0

#### Table 2.Distribution of various pesticide groups among contaminated vegetable groups.

Pestici	%	% contribution of vegetables and fruits to each group											
de	Contribut	Bri	Cabb	Okr	Tom	Green	Capsi	Caluli	Gra	Pom	Ban	Sweet	guava
group	ion	n	age	а	ato	chilli	cum	flower	pes	а	ana	orange	
		jal								gara			
										nte			
00	3.33	1.6	1.66	0	0	0	0	0	0	0	0	0	0
		6											
OP	35.0	3.3	3.33	6.6	8.33	8.33	1.66	1.66	3.33	1.66	0	0	0
		3		6									
SP	1.66	0	0	0	1.33	0	0	0	0	0	0	0	0

#### Table 3. Pesticide residue levsels in vegetables and fruit samples.

Pesticide	Variations in residue level	Mean <sup>a</sup> residues in	Standard deviation of
group	(ppm)	positive samples	residues in positive samples
			(ppm)
All pesticides	0.03-1.18	0.192	0.121
OC	0.087-0.10	0.071	0
OP	0.03-1.18	0.184	0.134
SP	0.88	0.88	0

#### Table 4.Concentration of OCs, OPs and SPs in Vegetable samples.

Name of the pesticides	% samples	Variations in residue level (ppm)	mean <sup>a</sup> (ppm)	SD <sup>b</sup> (ppm)
Organochlorines				
α - Endosulfan	1.66	0.10	0.1	0
β- Endosulfan	1.66	0.087	0.087	0
Organo				
Phosphates				
Dimethoate	13.3	0.05-1.18	0.309	0.371
Profenophos	6.66	0.08-0.59	0.212	0.217
Malathion	11.66	0.05-0.53	0.191	0.156
Chlorpyrifos	3.33	0.08-0.35	0.219	0.139
Quinalphos	6.66	0.03-0.09	0.075	0.026
Methyl	1.66	0.04	0.04	0
parathion				
Ethion	1.66	0.05	0.05	0

## ACKNOWLEDGEMENTS

The authors are Thankful to the Head of the Department of AINP on PRS, Acharya N.G Ranga Agricultural University, Hyderabad for his keen interest and providing research facilities.

## REFERENCES

- Adeyeye, A.&Osibanjo, O, 1999.Residues of organochlorine pesticides in fruits, vegetables and tubers from Nigerian markets. *Science of the Total Environment*, 231, 227–233.
- Agnihotri, N. P.1999. Pesticide, safety evaluation and monitoring. All India co-ordinated research project on pesticide residues pp. 119–146. New Delhi: Division of Agricultural Chemicals, *Indian Agricultural Research Institute*.
- Amoah, P., Drechsel, P., Abaidoo, R. C., and Ntow, W. J. 2006. Pesticide and pathogen contamination of vegetables in Ghana's urban markets. *Archives of Environmental Contamination and Toxicology*, 50, 1–6.
- Anastassiades, M. Lehotay, S.J., Stajnbaher, D., and Schenck, F. J. 2003. Fast and easy multiresidue method employing acetonitrile extraction/partitioning and "dispersive solidphase extraction" for the determination of pesticide residues in produce. *The Journal of AOAC International*, 86,412–431.
- Andaman and Nicobar Administration. 2009. Statistics book, Directorate of statistics. Port Blair: Andaman and Nicobar Administration.
- Bempah, C. K. Kwofie, A. B., Enimil, E., Blewu, B., and Martey, G. A. 2012. Residues of organochlorine pesticides in vegetables marketed in Greater Accra region of Ghana. *Food Control*, 25, 537–542.
- Bhanti, M. and Taneja, A. 2005. Monitoring of organochlorine pesticide residues in summer and winter vegetables from Agra, India—a case study. *Environmental Monitoring and Assessment*, 10, 341–346.
- Bhanti, M. and Taneja, A. 2007. Contamination of vegetables of different seasons with organophosphorus pesticides and related health risk assessment in northern India. *Chemosphere*, 69, 63–68.
- Bhattacharyya, A. Barik, S. R., and Ganguly, P. 2009. New pesticide molecules, formulation technology and uses: present status and future challenges. *The Journal of Plant Protection Sciences*, 11, 9–15.
- Charan, P. D. Ali, S. F., Yati, K., and Sharma, K. C.2010. Monitoring of pesticide residues in farmgate vegetables of central Aravalli region of western India. *American-Eurasian Journal of Agricultural & Environmental Science*, 73, 255–258.
- Chen, C. Yongzhong, Q., Qiong, C., Chuanjiang, T., Chuanyong, L., and Yun, L. 2011. Evaluation of pesticide residues in fruits and vegetable from Xiamen, China. *Food Control*, 227, 1114–1120.
- Chowdhury, M. T. I. Razzaque, M. A., and Khan, M. S. I.2011. Chlorinated pesticide residue status in tomato, potato and carrot. *Journal of Experimental Sciences*, 21, 1–5.

- Hjorth, K.Johansen, K., Holen, B., Andersson, A., Christensen, H. B., Siivinen, K., and Toome, M.2011.Pesticide residues in fruits and vegetables from South America—a Nordic project. *Food Control*, 2211, 1701–1706.
- Igbedioh, S. O.1991. Effects of agricultural pesticides on humans, animals and higher plants in developing countries. *Archives of Environmental Health*, 46, 218.
- Kalra, R. L. and Chawla, R. P.1985.Pesticidal contamination in foods in the year 2000 AD. Proceedings of *Indian National Science Academy*, B52, 188–204.
- Kumari, B. Kumar, R., Madan, V. K., Singh, R., Singh, J., and Kathpal, T. S. 2003. Monitoring ofpesticidal contamination in winter vegetables from Hisar, Haryana. *Environmental Monitoring and Assessment*, 87, 311–318.
- Kumari, B. Madan, V. K., Singh, J., Singh, S., and Kathpal, T. S.2004. Monitoring of pesticidal contamination of farmgate vegetables from Hisar. *Environmental Monitoring and Assessment*, 90, 65–71.
- Lozowicka, B. Jankowska, M., and Kaczynski, P. 2012. Pesticide residues in Brassica vegetables and exposure assessment of consumers. *Food Control*, 25, 561–575.
- Mandal, K. and Singh, B. 2010. Magnitude and frequency of pesticide residues in farmgate samples of cauliflower in Punjab, India. *Bulletin of Environmental Contamination and Toxicology*, 854, 423–426.
- Mathur, S.C., 1999. Future of Indian pesticides industry in next millennium. Pesticide Information, XXIV, 4, 9–23
- Mukherjee,I. 2003. Pesticides residues in vegetables in and around Delhi. Environmental *Monitoring and Assessment*, 86, 265–271.
- Nishina, T. Kien, C. N., Noi, N. V., Ngoc, H. M., Kim, C. S., Tanaka, S., and Iwasaki, K. 2010. Pesticide residues in soils, sediments and vegetables in the Red River Delta, northern Vietnam. *Environmental Monitoring and Assessment*, 169, 285–297.
- PFA. 1954. Prevention of Food Adulteration Act, 1954. Act. No. 37 with Prevention of Food Adulteration Rules, 1955 and Notification and Commodity Index. Lucknow: Eastern Book.
- Singh, D.P. 2001. Pesticides pollution on veterinary public health and food safety in India. In: Livestock Community and Environment. Proceedings of the 10th Conference of the Association of Institutions for Tropical Veterinary Medicine, Copenhagen, Denmark
- Singh, S. P. and Kiran Singh, N. K. 2006. Pesticideresidues in farmgate vegetable samples in Bihar. *Pest Management in Horticultural Ecosystems*, 122, 152–155.
- Waliszevski, S. M. PardioSedus, V. T., and Waliszevski, K. N. 1996. Detection of some organochlorine pesticides in cow's milk. *Food Additives and Contaminants*, 13, 231–235.

**Corresponding author: Dr. K. Lakshmi Devi**, Department of Biochemistry, S.K. University, Anantapuramu, Andhra Pradesh, India.

Email: <u>harinathreddya@gmail.com</u> <u>klakshmidevi2000@yahoo.co.in</u>