Genetic Diversity Based on Multivariate Analyses for Yield and it's Contributing Characters in Bread Wheat (*Triticum aestivum* L.) Genotypes

By

Birhanu Mecha, Sentayehu Alamerew, Alemayehu Assefa, Ermias Assefa and Dargicho Dutamo

ISSN 2319-3077 Online/Electronic ISSN 0970-4973 Print

UGC Approved Journal No. 62923 MCI Validated Journal Index Copernicus International Value IC Value of Journal 46.52 Poland, Europe (2015) Journal Impact Factor: 4.275 Global Impact factor of Journal: 0.876 Scientific Journals Impact Factor: 3.285 InfoBase Impact Factor: 3.66

J. Biol. Chem. Research Volume 34 (2) 2017 Pages No. 440-453

Journal of Biological and Chemical Research

An International Peer Reviewed / Referred Journal of Life Sciences and Chemistry

Indexed, Abstracted and Cited in various International and National Scientific Databases

Published by Society for Advancement of Sciences®

Received: 06/07/2017



RESEARCH PAPER Accepted: 22/09/2017

Genetic Diversity Based on Multivariate Analyses for Yield and it's Contributing Characters in Bread Wheat (*Triticum aestivum* L.) Genotypes

Revised: 20/09/2017

Birhanu Mecha, *Sentayehu Alamerew, **Alemayehu Assefa, ***Ermias Assefa and ****Dargicho Dutamo Wachemo University, College of Agricultural Sciences, Hosanna, Ethiopia *Jimma University, College of Agriculture and Veterinary Medicine, Ethiopia. **Ethiopian Institute of Agricultural Research (EIAR); Addis Ababa, Ethiopia. ***Ethiopian Biotechnology Research Institute, Ethiopia,(EBTI). ***Mizan-Tepi University, college of Agriculture and natural resources, Mizan Teferi, Ethiopia.

ABSTRACT

Breed wheat is one of the most important export and strategic cereal crop in the world and in Ethiopia in terms of production and utilization. Information about genetic diversity is important for selection of promising parents in hybrid variety and inbred line development, in assigning inbred lines to heterotic groups, variety registration and protection and to estimate the potential of genetic gain in a breeding program. The overall objective was to assess the genetic diversity of wheat genotypes by multivariate analyses (Cluster and Principal component). The experimental material consisted of sixty four breed wheat genotypes tested in 8x8 Simple Lattice Design at Ginchi. Analysis of variance revealed highly significant differences among accessions for all traits. Cluster analysis revealed that the 61 genotypes and 3 checks were grouped into eight clusters. The crosses between genotypes selected from cluster-V with cluster-VIII and cluster V with cluster VII are expected to produce better genetic recombination and segregation in their progenies. Five principal components (PC1, PC2, PC3, PC4 and PC5) exhibited eigen value greater than one and accounted 72.78% of the total variation.

Keywords: Cluster, Principal Component, Genetic Recombination and Segregation.

INTRODUCTION

Bread wheat (*Triticum aestivum* L. em Thell. 2n=6x=42), a self-pollinating annual plant in the true grass family *Gramineae* (*Poaceae*), is the largest cereal crop extensively grown as staple food source in the world (Mollasadeghi and Shahryari, 2011).

Cluster analysis is a multivariate method, which aims to classify a sample of subjects based on a set of measured variables into a number of different groups such that similar subjects are placed in the same group. It sorts genotypes into groups, or clusters, so the degree of association will be strong between members of the same cluster and weak between members of different clusters. The cluster analysis was performed using a measure of similarity levels and Euclidean distance (Everitt, 1993; Eisen *et al.*, 1998).

Different researcher grouped bread wheat genotypes using cluster analysis. Hailegiorgis *et al.* (2011) reported that the cluster analysis grouped the 49 bread wheat genotypes into 22 different clusters. This indicates the presence of wide diversity among the tested genotypes. Khodadadi *et al.* (2011) determined the genetic diversity of 36 winter wheat cultivars from Iran and by using cluster analysis, seven clusters were determined.

Principal component analysis makes it possible to transform a given set of characteristics (variables), which are mutually correlated, into a new system of characteristics, known as principal components, which are not correlated. The obtained variables may also be used for further analysis, where the assumption of no co-linearity is required. Moreover, the analysis is characterized by the fact that it includes the total variance of variables, explains maximum of variance within a data set, and is a function of primary variables (Katarzyna *et al.* 2012).

Precise knowledge about germplasm diversity and genetic relationship among breeding materials is a pre-requisite for crop improvement programs as it helps in the development of superior recombinants (Rauf *et al.* 2012). Genetic divergence analysis estimates the extent of diversity existed among selected genotypes (Mondal and Khajuria, 2003). Precise information on the nature and degree of genetic diversity helps the plant breeder in choosing the diverse parents for purposeful hybridization (Hailegiorgis *et al.*, 2011). Generating fertile genetic diversity information among wheat genotypes is very important tool because the information will help the wheat breeders to bread for many characters (earliness, yield increase, drought tolerance, etc). Within the investigated material and this new material will serve as a new stock for the improvement of wheat breeding program for traits of interest. Therefore, the objective of this study was to assess the genetic diversity of durum wheat accessions by multivariate analyses.

MATERIALS AND METHODS

Description of the study area

The experiment was conducted at Ginchi, West Shewa in 2012/13 cropping season. Ginchi Agricultural Research Sub Center is located at an altitude of 2240 meters above sea level, 84 kilometers (kms) to the West of Addis Ababa, and at a Latitude and Longitude of 09°03'N and 38°15'E, respectively. It is the center where the cereal crops like Teff, barley and wheat are grown. The maximum and minimum temperatures of the area are 24.72 °C and 8.76 °C, respectively, whereas the mean annual rainfall is 1080.4mm. The major soil types are black (Vertisol) and clay loam with pH of 6.4, which is heavy clay with 0.91-1.32% organic matter (HARC, Soil Analysis and Plant Physiology Team, 2012).

Experimental Material

A total of sixty four bread wheat (*Triticum aestivum L*.) genotypes that include three standard checks and sixty one exotic bread wheat accessions introduced from CIMMYT were included in this study. The accessions were obtained kindly from HARC. The three released cultivars Digelu, Alidoro and Meraro were used as a standard checks. They were selected based on their agronomic performances and suitability to the growing conditions (Table 1).

Experimental Design and Trial Management

The experiment was carried out in 8x8 Simple Lattice Design at random. The genotypes were grown under uniform rain fed conditions. The plot size was six rows of 2.5 m length with 0.2 m row spacing i.e. $1.2 \text{ m x } 2.5 \text{ m} = 3\text{m}^2$ (standard plot size for variety trial).

Planting was done by hand drilling on July 06, 2012. Seed rate was 150 kg/ha (45 g/plot). Recommended fertilizer rate of 100/100 kg/ha N/P_2O_5 in the forms of Urea and DAP was applied to each plot in the shallow furrow depths and mixed with soil at the same time during sowing. For data collection, the middle four rows were used ($2m^2$ area). The central four rows were harvested for grain yield and biomass yield from each plot leaving boarder rows to avoid boarder effects. All other agronomic practices were undertaken uniformly to the entire plot as recommended for wheat production in the area during the growing season.

Entry	Pedigree	Seed source	Entry	Pedigree	Seed source
1	CIMMYTOB/2	CIMMYT	33	CIMMYTOB/65	CIMMYT
2	CIMMYTOB/7	CIMMYT	34	CIMMYTOB/66	CIMMYT
3	CIMMYTOB/14	CIMMYT	35	CIMMYTOB/67	CIMMYT
4	CIMMYTOB/22	CIMMYT	36	CIMMYTOB/68	CIMMYT
5	CIMMYTOB/23	CIMMYT	37	CIMMYTOB/70	CIMMYT
6	CIMMYTOB/24	CIMMYT	38	CIMMYTOB/71	CIMMYT
7	CIMMYTOB/25	CIMMYT	39	CIMMYTOB/75	CIMMYT
8	CIMMYTOB/27	CIMMYT	40	CIMMYTOB/76	CIMMYT
9	CIMMYTOB/29	CIMMYT	41	CIMMYTOB/77	CIMMYT
10	CIMMYTOB/32	CIMMYT	42	CIMMYTOB/78	CIMMYT
11	CIMMYTOB/33	CIMMYT	43	CIMMYTOB/79	CIMMYT
12	CIMMYTOB/35	CIMMYT	44	CIMMYTOB/80	CIMMYT
13	CIMMYTOB/39	CIMMYT	45	CIMMYTADT/1	CIMMYT
14	CIMMYTOB/40	CIMMYT	46	CIMMYTADT/2	CIMMYT
15	CIMMYTOB/41	CIMMYT	47	CIMMYTADT/3	CIMMYT
16	CIMMYTOB/44	CIMMYT	48	CIMMYTADT/4	CIMMYT
17	CIMMYTOB/45	CIMMYT	49	CIMMYTADT/5	CIMMYT
18	CIMMYTOB/48	CIMMYT	50	CIMMYTADT/6	CIMMYT
19	CIMMYTOB/49	CIMMYT	51	CIMMYTADT/7	CIMMYT
20	CIMMYTOB/50	CIMMYT	52	CIMMYTADT/8	CIMMYT
21	CIMMYTOB/51	CIMMYT	53	CIMMYTADT/9	CIMMYT
22	CIMMYTOB/52	CIMMYT	54	CIMMYTADT/11	CIMMYT
23	CIMMYTOB/53	CIMMYT	55	CIMMYTADT/13	CIMMYT
24	CIMMYTOB/54	CIMMYT	56	CIMMYTADT/15	CIMMYT
25	CIMMYTOB/57	CIMMYT	57	CIMMYTADT/16	CIMMYT
26	CIMMYTOB/58	CIMMYT	58	CIMMYTADT/17	CIMMYT
27	CIMMYTOB/59	CIMMYT	59	CIMMYTADT/19	CIMMYT
28	CIMMYTOB/60	CIMMYT	60	CIMMYTADT/20	CIMMYT
29	CIMMYTOB/61	CIMMYT	61	CIMMYTADT/21	CIMMYT
30	CIMMYTOB/62	CIMMYT	62	ALIDORO	HARC
31	CIMMYTOB/63	CIMMYT	63	MERARO	KARC
32	CIMMYTOB/64	CIMMYT	64	DIGELU	KARC

Table 1. List of genotypes used in the study.

Description of Data Collected

The data on the following attributes was collected on the basis of the central four rows in each plot per replication.

Days to 50% heading (DH): The numbers of days from sowing to 50% of plants have started heading. **Days to 75% maturity (DM):** The numbers of days from date of sowing to a stage at which 75% of the plants have reached physiological maturity or 75% of the spikes on the plots turned golden yellow color.

Grain filling period: The grain filling period in days was computed by subtracting the number of days to heading from the number of days to maturity.

Thousand Kernels weight (TKW) (g): The weight of 1000 kernels from randomly sampled seeds per plot measured with sensitive balance.

Grain yield per plot (GYP) (g): The grain yield per plot was measured using sensitive balance after moisture of the seed is adjusted to 12.5%. Total dry weight of grains harvested from the middle four rows was taken as grain yieldt.

Biomass yield per plot (BMYP): It was recorded by weighing the total above ground yield harvested from the four central rows of each experimental plot at the time of harvest.

Harvest index (%): It was estimated by dividing grain yield per plot to biological yield per plot.

Hectoliter weight (HLW) (kg/ha): It is grain weight of one hectoliter volume random sample of wheat grain for each experimental plot.

Ten plants were randomly selected from the four central plots for recording the following observations:

Plant height (cm): The average height of ten randomly taken plants at the maturity time from the middle four rows of each plot of the replication was measured from the ground level to the top of the spike excluding the awn.

Number of productive tillers per plant: The numbers of tillers per plant bearing productive heads were counted at the time of harvest and average was recorded for the ten randomly taken plants from the middle four rows.

Spike length (cm): The average spike length of ten randomly taken plants from the base of the main spike to the top of the last spikelet excluding awns was recorded from four central rows of each plot. **Number of spikelets per spike:** Total number of spikelets on main spike of all ten plants from four central rows were counted at the time of maturity and average was recorded.

Number of kernels per spike (NKPS): Total number of grains in the main spike were counted at the time of harvest from ten randomly taken plants and expressed as average and recorded from four central rows of each plot.

Statistical Analysis

Analysis of variance (ANOVA)

The data collected for each quantitative trait were subjected to analysis of variance (ANOVA) for simple lattice design. Analysis of variance was done using Proc lattice and Proc GLM procedures of SAS version 9.2 (SAS Institute, 2008) after testing the ANOVA assumptions.

The mathematical model for Simple Lattice Design is:

$Y_{ijr} = \mu + A_r + G_{ij} + B_{ir} + B_{jr} + e_{ijr},$

where Y_{ijr} = the value observed for the plot in the rth replication containing the genotype G_{ij} , μ = grand mean, G_{ij} = genotype effect in the ith row and jth column, A_r = replication effect, B_{ir} =ith block effect, B_{jr} =jth block effect, e_{ijr} , = the plot residual effect.

Cluster analysis

Clustering the genotypes into different groups was carried out by average linkage method. The appropriate number of clusters was determined from the values of Pseudo F and Pseudo T^2 statistics using the procedures of SAS computer software version 9.2 to group sets of genotypes into homogeneous clusters (SAS Institute, 2008).

Genetic divergence analysis

Genetic divergence between clusters was determined using the generalized Mahalanobis D^2 statistics (Mahalanobis, 1936).

The D^2 analysis performed based on the mean values of all traits by using SAS software program. In matrix notation, the distance between any two groups was estimated from the following relationship.

$$D^{2}_{ij} = (Xi - Xj) S^{-1} (Xi - Xj)$$

Where, D_{ij}^2 = the square distance between any two accessions i and j; X_i and X_j = the vectors for the values for accession ith and jth genotypes; and S⁻¹ = the inverse of pooled variance covariance matrix within groups.

Testing the significance of the squared distance values obtained for a pair of clusters was taken as the calculated value of χ^2 (chi-square) and tested against the tabulated χ^2 values at n-2 degree of freedom at 1% and 5% probability level, where n= number of characters used for clustering genotypes.

Principal component analysis (PCA)

The principal component analysis was performed using the proc princomp procedure of SAS version 9.2 software (SAS, 2008). Statistical inference was computed by taking into account all the factors at a time. In this study, investigation of suitable multivariate technique for analyzing data for all the characters are proposed. The general formula to compute scores on the first component extracted in a principal component analysis:

$$PC1 = b11(X1) + b12 + \cdots b1p = (Xp)$$

where, PC1 = the subject's score on principal component 1 (the first component extracted), b1p = the regression coefficient (or weight) for observed variable p, as used in creating principal component 1 and Xp = the subject's score on observed variable p.

RESULTS AND DISCUSSION

Analysis of variance (ANOVA)

Mean squares of the 13 traits from analysis of variance (ANOVA) are presented in Table 2. Highly significant differences among genotypes (P<0.01) were observed for seven characters (days to heading, number of productive tillers per plant, spike length, number of spikelets per spike, 1000 kernel weight, grain yield plot⁻¹ and hectoliter weight or test weight), significant at (P<0.05) for the rest six characters; namely, days to 75% maturity, grain filling period, plant height, number of kernels spike⁻¹, biomass yield and harvest index. This result indicates that there is variability among the studied genotypes.

Genetic divergence

Genetic divergence analysis quantifies the genetic distance among the selected genotypes and reflects the relative contribution of specific traits towards the total divergence. The genetic improvement through hybridization and selection depends upon the extent of genetic diversity between parents (Hailu *et al*, 2016)

Clustering of genotypes

The dendrogram obtained from the cluster analysis grouped the 64 bread wheat genotypes in to seven groups and one solitary (Fig 1). The F-test for the seven clusters indicated that there were statistically accepted differences between clusters. The genotypes were clustered in such a way that 8 genotypes (12.5%) were grouped into cluster I, 9 genotypes (14.0625%) into cluster II, 11

genotypes (17.1875%) into cluster III, 5 genotypes (7.8125%) and 9 genotypes (14.0625%) into clusters IV and V, respectively, 18 genotypes (28.125%) into cluster VI, 3 genotypes (4.6875%) into cluster VII as well as one genotype (1.5625) into cluster VIII Table 3. Similar works were done by Shashikala (2006) grouped 169 wheat genotypes in to 11 clusters, Noorka and Khaliq (2007) grouped 100 bread wheat genotypes in to 13 clusters; Degewione and Alamerew (2013) grouped 26 bread wheat genotypes into six clusters.

		genotypes.			
Characters	Replication	Genotype	Intra Block	CV (%)	Efficiency
	(df=1)	(df=63)	Error		Relative to
			(df=49)		RCBD
Days to 50% heading	8.51	86.99**	9.22	10.62	107.38
(days)					
Days to 75% maturity	2.53	37.75*	5.42	4.04	100.49
(days)					
Grain filling period (days)	13.78	47.30*	5.58	10.16	100.72
Plant height (cm)	29.55	248.40*	22.48	11.94	101.24
Number of productive	0.02	0.68**	0.16	10.58	105.37
tillers per plant					
Spike length (cm)	0.71	1.09**	0.10	8.84	109.36
Number of spikelets per	3.30	2.54**	0.25	6.87	102.31
spike					
Number of kernels per	25.92	50.45*	8.83	16.20	111.15
spike					
1000 kernels weight (g)	0.13	43.21**	3.72	12.63	103.52
Biomass yield per plot (g)	22578	160197*	9604	19.50	115.26
Harvest index (%)	18.90	37.24*	6.19	18.09	103.12
Hectoliter weight (kg/hL)	6.71	18.43**	3.42	13.75	116.27
Grain yield per plot (g)	1287.78	22864**	4066	2.07	120.90

Table 2. Analysis of variance (mean squares) for the 13 characters of analyzed bread wheat

Df =Degrees of freedom

*=significant at 5% probability level and **=highly significant at 1% probability level CV= Coefficient of Variation, RCBD=Randomized Complete Block Design

Cluster mean analysis

The mean values of the thirteen traits in each cluster are presented in Table 4. Cluster I exhibited the highest days to maturity and the delayed maturity time (127.5 days) with largest number of kernels per spike (43.89); thus containing late maturing genotypes. Cluster II could be characterized by the earliest maturity time (114.890 days) and low number of spikelets per spike (16.95) but with the highest harvest index (40.1%). Cluster III had the lowest number of productive tillers per plant (5.736 with the lowest biomass yield per plot (1654.6g).

A relatively larger range between clusters was displayed for plant height. Cluster V had the shortest plant height (90.28cm) and cluster IV revealed the highest (122.15cm) with the highest number of spikelets per spike (18.45), but it showed the lowest harvest index (31.2%). The highest number of days to fill grain was observed in cluster VIII (63.00 days) and the lowest goes for cluster I (52.875 days). The ranges of number of spikelets spike⁻¹, thousand kernels weight (g), harvest index (%), spike length (cm) and hectoliter weight (kg/hL) were low.



Figure 1. Dendrogram showing the clusters of sixty four bread wheat genotypes.

Cluster	Number of Genotypes	Accession Code
I	8	ALIDORO,CIMMYTOB/23,CIMMYTOB/40,CIMMYTADT/1, CIMMYTADT/13,
		CIMMYTOB/45,CIMMYTOB/49,CIMMYTOB/50,CIMMYTOB/51,
11	9	CIMMYTOB/52,CIMMYTOB/60,CIMMYTOB/61, IMMYTOB/62,
		CIMMYTADT/2
		CIMMYTOB/54,CIMMYTOB/57,CIMMYTOB/58,CIMMYTOB/63,
III	11	CIMMYTOB/76,CIMMYTADT/3,CIMMYTADT/4,CIMMYTADT/5,
		CIMMYTADT/8, CIMMYTADT/11, CIMMYTADT/20
IV	5	CIMMYTOB/14,CIMMYTOB/33,CIMMYTOB/35,CIMMYTADT/6, CIMMYTADT/7
M	9	CIMMYTOB/22,CIMMYTOB/27,CIMMYTOB/32, CIMMYTOB/39,
v		CIMMYTOB/41,CIMMYTOB/48,CIMMYTADT/9,CIMMYTADT/21, MERARO
		CIMMYTOB/2,CIMMYTOB/22,CIMMYTOB/53,CIMMYTOB/59,
		CIMMYTOB/64,CIMMYTOB/65,CIMMYTOB/66,CIMMYTOB/67,
VI	18	CIMMYTOB/68, CIMMYTOB/70, CIMMYTOB/71, CIMMYTOB/75,
		CIMMYTOB/77,CIMMYTOB/78,CIMMYTOB/79,CIMMYTOB/80,
		CIMMYTADT/15, CIMMYTADT/19,
VII	3	CIMMYTOB/24, CIMMYTOB/29, CIMMYTOB/44
VIII	1	CIMMYTOB/25

Table 3. Distribution of bread wheat genotypes in different cluster groups.

Cluster V showed the shortest plant height (90.28cm), the lowest yield per plot (642.8g) and the lowest hectoliter weight (77.87 kg/hL) but revealed the latest days to heading (74.667 days) and maturity (126.778 days).

J. Biol. Chem. Research

Cluster VI showed the longest spike length (8.92cm) and the highest hectoliter weight (80.79 kg/hL). Cluster VII revealed intermediate characteristics for all of the traits. Cluster VIII revealed the highest grain yield (1182g), the heaviest thousand kernels weight (48.20g) and the highest biological yield per plot (3500g) but the lowest number of kernels per spike (24.00).

Relative contribution of each character towards divergence

The analysis of the contribution of each character towards the expression of genetic divergence (Table 5) indicated that days to 50% heading contributed maximum followed by days to 75% maturity to the total genetic divergence in the genotypes studied. These two traits followed by grain filling period, plant height, number of productive tillers per plant, days to 75% maturity, spike length, number of spikelets per spike, number of kernels per spike and thousand kernels weight totally accounted for 94.65% of total genetic divergence in the materials studied. Grain yield, biomass yield, harvest index and hectoliter weight were the least contributor to the divergence.



Figure 2. Constellation plot showing the clusters of sixty four bread wheat genotypes.

Traits /	I	II		IV	V	VI	VII	VIII
Clusters								
Days to	74.625	60.333	62.909	70.600	74.67**	62.056	59.667	59.00*
heading								
(days)								
Days to 75%	127.50**	114.89*	116.818	124.800	126.778	123.389	121.000	122.000
maturity								
(days)								
Grain filling	52.875*	53.444	53.181	54.200	52.889	60.944	61.333	63.00**
period(days)								
Plant	108.375	92.720	92.091	122.15**	90.28*	92.2361	119.250	115.750
height(cm)								
No. of	6.207	5.994	5.74*	6.646	6.767	6.119	6.417	6.80**
productive								
tillers plant ⁻¹								
Spike	8.550	8.661	8.600	8.430	8.483	8.92**	8.600	8.25*
length(cm)								
Number of	18.132	16.95*	17.295	18.45**	18.361	17.044	17.517	18.050
spikelets								
spike ⁻¹								
Number of	43.89**	40.544	31.181	34.960	39.467	38.322	41.633	24.00*
kernels spike								
1								
Thousand	36.013*	40.822	36.39	35.040	30.844	42.950	43.167	48.20**
kernels								
weight(g)								
Grain yield	756.8	825.4	643	768.6	642.8*	768.6	857.4	1182**
plot⁻¹(g)								
Biomass yield	2092.6	2100	1655*	2360	2066	2061	2333.4	3500**
plot⁻¹(g)								
Harvest index	34.4	40.1**	39.7	31.2*	31.6	37.2	36.7	34.0
(%)								
Hectoliter	80.388	79.567	79.936	79.840	77.87*	80.79**	80.933	80.50
weight(kg/hL)								

Table 4.Mean values of 13 characters of the eight clusters for the 64 bread wheat genotypes.

* =Lowest value, **= Highest value.

Average intra and inter cluster distance (D²)

The average intra and inter cluster distance D^2 values are presented in Table 6. Maximum average intra cluster D^2 was shown by cluster III (24.35) followed by cluster VI (23.99). The lowest intra cluster distance D^2 was recorded in cluster VII (9.40), which shows the presence of less genetic variability or diversity within this cluster. The diversity among clusters or inter cluster distance D^2 ranged from 85.15 to 174.32. Cluster V and VIII showed maximum inter cluster distance of 174.32, followed by that between clusters V and VII (161.64). The lowest inter cluster distance was noticed between clusters II and III (85.15), followed by that between clusters II and VI (88.77). Evaluation of genetic diversity can be useful for the selection of the most efficient genotypes.

		alversity.
Character	Rank	Contribution (%)
Days to 50% heading	520	25.81
Days to 75% maturity	382	18.93
Grain filling period	220	10.95
Plant height	184	9.14
Number of productive tillers per plant	160	7.95
Spike length	132	6.55
No. of spikelets per spike	109	5.41
No. of kernels per spike	105	5.19
1000-kernels weight	96	4.74
Grain yield	52	2.6
Biomass yield	45	2.24
Harvest index	6	0.28
Hectoliter weight	5	0.23

Table 5. Contribution of each trait towards total diversity.



Rank

Fig. 3 Contribution of each trait towards total diversity.

Table 6. Average intra (bold) and inter cluster (off diagonal) D^2 v	values among eight clusters in
bread wheat genotypes.	

Clusters	I		III	IV	V	VI	VII	VIII
	12.20	105.45**	96.16**	102.26**	99.53**	144.11**	142.68**	128.11**
=		10.88	85.15*	157.76**	129.46**	87.95*	118.54**	140.51**
=			24.35	152.54**	88.77*	94.27**	156.59**	142.37**
IV				12.00	152.90**	155.87**	107.5**	94.34**
V					23.86	112.98**	161.64**	174.32**
VI						23.99	117.47**	99.31**
VII							9.40	89.75*
VIII								0

 x^2 =82.529 at 5% probability level and x^2 =92.010 at 1% probability level, *= Significant at 0.05 probability level, **= Highly significant at 0.01 probability level

The results of this study showed the presence of a high genetic divergence among wheat genotypes, similar to the findings of Ali et al. (2008) who reported that cluster analysis can be useful for finding high yielding wheat genotypes.

According to Rahim *et al.* (2010) who showed that the hybrids of genotypes with maximum distance resulted in high yield, the cross between these genotypes can be used in breeding programs to achieve maximum heterosis. Therefore, more emphasis should be given on cluster V and VIII for selecting genotypes as parents for crossing with the genotypes of cluster, which may produce new recombinants with desired traits. The chi-square test for the clusters indicated that there was a statistically significant difference in all characters (Table 7). The χ^2 - test for the eight clusters indicated that there was a statistically significant difference in all characters.



Fig.4 Average intra and inter cluster distance.

Principal component analysis

Principal component analysis reflects the importance of the largest contributor to the total variation at each axis for differentiation (Sharma, 1998).The data matrix of 13*64 was prepared for principal component analysis. Out of thirteen principal components (PCs), the first five exhibited eigenvalue greater than one (significant). The rest eight PCs explained non-significant amount of variation. The eigenvalues are used to determine how many factors to retain. The sum of the eigen values is usually equal to the number of variables. Therefore, in this analysis the first factor retains the information contained in 3.36 of the original variables. The coefficients defining the first five principal components of these data are given in Table 7. The principal component analysis revealed that five principal components (PC1-C5) exhibited eigen value higher than one, with values 3.36, 2.46, 1.43, 1.19 and 1.03, respectively, have accounted for 72.78% of the total variation so these five were given due importance for further explanation.

According to Chahal and Gosal (2002), characters with largest absolute value closer to unity within the first principal component influence the clustering more than those with lower absolute value closer to zero. Therefore, in the present study, differentiation of the genotypes into different clusters was because of relatively high contribution of few characters rather than small contribution from each character. Accordingly, the first principal component (PC1) which accounted for 25.81% of the variability among genotypes were attributed to discriminatory traits such as days to heading, days to maturity, grain filling period, plant height, number of spikelets per spike, thousand kernels weight, harvest index and hectoliter weight. Likewise, 18.93% of the total variability among the tested genotypes accounted for the second principal component (PC2) mainly extracted from variation in days to maturity, grain filling period, plant height, number of productive tillers per plant, number of kernels per spike, grain yield per plot, biomass yield and test weight.

Similarly, the major contributing characters for the 10.94% of total variation in the third principal component (PC3) were spike length, grain filling period, grain yield, harvest index, thousand kernels weight, and days to maturity and number of productive tillers per plant. Furthermore, the fourth principal component (PC4), which explained 9.14% of total variations, was obtained from days to maturity, spike length, number of spikelets per spike and hectoliter weight. Quantitative characters such as plant height, number of kernels per spike, biomass yield per plot and harvest index, explained mainly for the remaining 7.95% of the variations of fifth principal component (PC3) (Table 8).

				.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
Characters		PC1	PC2	PC3	PC4	PC5
Days to 50% heading		0.907	-0.061	-0.060	-0.101	0.161
Days to 75% maturity		0.613	0.371	0.362	-0.355	0.243
Grain filling period		-0.516	0.444	0.436	-0.277	0.034
Plant height		0.325	0.539	0.080	-0.205	-0.318
Number of productive tillers per pla	nt	0.279	0.324	-0.321	-0.291	0.093
Spike length		0.153	0.160	0.623	0.607	0.117
Number of spikelets per spike		0.747	0.123	0.132	0.370	-0.005
Number of kernels per spike		0.208	0.438	-0.071	0.166	0.634
Thousand kernels weight		-0.679	0.277	0.390	0.148	-0.124
Grain yield per ha		-0.195	0.796	-0.423	0.282	-0.064
Biomass yield per ha		0.193	0.799	-0.189	0.114	-0.391
Harvest index		-0.578	0.101	-0.422	0.229	0.404
Hectoliter weight		-0.459	0.368	0.182	-0.384	0.292
Eigen value	3.36	2.46	5 1.43	1.19	1.03	
Difference	0.90	1.03	3 0.24	0.16	0.18	
Proportion	25.8	1 18.9	3 10.95	9.14	7.95	
Cumulative	25.82	1 44.7	4 55.69	64.83	72.78	

Table 7. Eigen values and Eigenvectors of the first five principal components (PCs) for 13
characters of bread wheat genotypes.



Fig 5. graphical presentation of the first five principal components.

SUMMARY AND CONCLUSION

Clustering was made to categorize quantitative traits into components for the sake of understanding the share components contribute to major variation in the study. The cluster analysis based on D^2 analysis on pooled mean of genotypes classified the 64 genotypes into eight clusters; which makes them to be moderately divergent. There was statistically approved difference between all the characters. Mean values in each cluster revealed that genotypes in cluster I had relatively moderate in most of characteristics but late heading and maturity days with the shortest time to fill the kernels. Mean values in genotypes grouped in cluster II had low to moderate characteristics with earliest maturity time and high number of kernels per spike with high thousand kernels weight. Mean values of cluster III showed the least number of productive tillers per plant, short time to fill the kernels, the earliest maturing types, low in grain and biomass yields and the lowest number of kernels per spike. The highest inter-cluster distance were exhibited between cluster V and VIII ($D^2 =$ 174.32), cluster V and VII (D^2 = 161.64) and cluster II and IV (D^2 = 157.76), indicates wider genetic divergence among the clusters. Whereas, the shortest squared distance was observed between cluster II and III ($D^2 = 85.15$) at, followed by between cluster II and VI ($D^2 = 88.77$). The crosses between genotypes selected from cluster-V with cluster-VIII ,cluster V with cluster VII and cluster II and IV are expected to produce better genetic recombination and segregation in their progenies. Therefore, these bread wheat genotypes need to be crossed and selected to develop high yielding variety. The principal component analysis revealed that principal components PC1, PC2, PC3, PC4 and PC5, accounted for 72.78% of the total variation. This result further confirmed the presence of ample genetic diversity for use in improvement program.

ACKNOWLEDGMENTS

The authors would like to thank Jimma University College of Agriculture and Veterinary Medicine (JUCAVM) and Holeta Agricultural Research Center (HARC) for their financial support.

REFERENCES

- Ali, Y., Atta, B.M., Akhter, J., Monneveux, P. and Lateef, Z. (2008). Genetic variability, association and diversity studies in wheat (Triticum aestivum L.) germplasm. Pak J Bot 40: 2087- 2097.
- **Chahal, G.S. and S.S. Gosal, 2002.** Principles and Procedures of Plant Breeding: Biotechnology and Conventional Approaches. Narosa Publishing House, New Delhi, India, pp: 604.
- Daniel Hailegiorgis, Mebrahtom Mesfin and Tsige Genet ,2011. Genetic Divergence Analysis on some Bread Wheat Genotypes Grown in Ethiopia, 2Bahir Dar University, Amhara, Ethiopia.
- **Degewione A. and Alamerew S., 2013.** Genetic Diversity in Bread Wheat (*Triticum aestivum* L.) Genotypes. Pakistan Journal of Biological Sciences, 16: 1330-1335.
- **Eisen MB, Spellman PT, Brown PO, Botstein D, 1998.** Cluster analysis and display of genome-wide expression ilatteriis. Proceedings of the National Academy of Science, USA 95: 14863–14868. Ethiopia. Doctoral (Agri.) Thesis University of Agricultural Sciences.Alnarp,Sweden.
- Everitt BS, 1993. Cluster Analysis. Wiley, New York, NY.
- Hailu, A., Alamerew, S., Nigussie, M., & Assefa, E. (2016). Study of Genetic Diversity in Different Genotypes of Barley (Hordeum vulgare L.) Based on Cluster and Principal Component Analyses, Agricultural Science Research Journal. 6(February), 31–42. http://resjournals.com/journals/agricultural-science-research-journal.html
- Katarzyna Rymuza, Elżbieta Turska, Grażyna Wielogórska, Antoni Bombik, 2012. Use of principal component analysis for the assessment of spring wheat characteristics. Acta Sci. Pol., Agricultural 1(1): 79-90.

Mahalanobis, P.C., 1936. On the generalized distance in statistics. Proc. Nat. Inst. Sci., 2: 49-55

Mollasadeghi, V., Shahryari, R. 2011. Important morphological markers for improvement of yield in bread wheat. Advances Environ. Biol., 5 (3): 538 – 542.

- Mondal, S.K. and Khajuria, M.R. 2001. Correlation and path analysis in bread wheat (*Triticum aestivum* L.) under rainfed condition. Environment and Ecology, 18 (2): 405 408.
- Mostafa Khodadadi, Mohammad Hossein Fotokian, and Mohammad Miransari, 2011. Genetic Diversity of wheat (*Triticum aestivum* L.) genotypes based on cluster and principal component analyses for breeding strategies, Australian Journal of Crop Sciences, Tehran, Iran.
- Noorka, I.R. and I. Khaliq, 2007. An efficient technique for screening wheat (*Triticum aestivum* L.) germplasm for drought tolerance. Pak. J. Bot., 39: 1539-1546.
- Rahim, M.A., Mia, A.A, Mahmud, F., Zeba, N. and Afrin, K. 2010. Genetic variability, character association and genetic divergence in murgbean platn Omic 3:1-6.
- Rauf, S., Tariq, S.A., Hassan, S.W. (2012). Estimation of pedigree based diversity in Pakistani wheat (*Triticum aestvium* L.) germplasm. Communications in Biometry and Crop Science 7 (1), 14– 22
- SAS, 2008. Statistical Analysis System. Version 9.2, SAS Institute Inc., Cary, NC., USA.
- Sharma, P.K., Gupta, P.K. and Balayn, H.S., 1998, Genetic diversity in a large collection of wheats (*Triticum* spp.). *Indian Journal of Genetics and Plant Breeding*, 58(3): 271-278.
- Shashikala, S.K. 2006. Analysis of genetic diversity in wheat. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Dharwad, India.

Corresponding author: Ermias Assefa, Jimma University, College of Agriculture and Veterinary Medicine, Ethiopia. Email: <u>ethioerm99@gmil.com</u>