

The Detoxification Effect of Nitrogen dose on Cadmium Toxicity in *Nicotiana tabacum*

By

Leila Saafi, Houda Gouia and Chiraz Chaffei Haouari


ISSN 0970-4973 (Print)

ISSN 2319-3077 (Online/Electronic)

Volume 29
No.2 (2012)

J. Biol. Chem. Research
Volume 29 2012 Pages No. 262-272

**Journal of
Biological and
Chemical Research**



Published by Society for Advancement of Sciences®

J. Biol. Chem. Research. Vol. 29, No. 2: 262-272 (2012)

(An International Journal of Life Sciences and Chemistry)

ms 29/2/24/2012, All rights reserved

ISSN 0970-4973 (Print)

ISSN 2319-3077 (Online/Electronic)

Published by Society for Advancement of Science®



JBCR

[http:// www.jbcr.in](http://www.jbcr.in)

jbiolchemres@gmail.com

info@jbcr.in

RESEARCH PAPER

Received: 01/08/2012 Revised : 03/10/2012 Accepted: 17/08/2012

The Detoxification Effect of Nitrogen dose on Cadmium Toxicity in *Nicotiana tabaccum*

Leila Saafi, Houda Gouia and Chiraz Chaffei Haouari

Unite de Recherche : Nutrition et Métabolisme Azotés et Protéines de Stress (99UR/09-20), Département de Biologie, Faculté des Sciences de Tunis. Campus Universitaire El Manar I, 1060, Tunis- Tunisie.

ABSTRACT

*In this study, researchers used *Nicotiana tabaccum*, as a model species to study its morphological, physiological and biochemical responses to the interaction of different concentrations of nitrogen and Cd, discussing the detoxification effect of nitrogen to Cd stress in *Nicotiana tabaccum*. The treatments were consisted of three Cd levels (0, 5 and 50 μ M) and three N levels (0, 0.1 and 5 mM). After 7 days treatment, the results indicated that at the same N level, Cd treatment had significant effect on all measured parameters, including plant growth, chlorophyll concentration, Malondialdehyde (MDA) concentration, antioxidant enzymes activities (SOD, POD, CAT and APX) and free proline. Meanwhile, study also showed there was significant difference among the three levels of N treatment. With the increasing concentration of N supplement, it showed more significant effect of alleviation to Cd induced damage in *Nicotiana tabaccum*.*

Key words: *Nicotiana tabaccum*, cadmium-N interaction, antioxidant enzymes

INTRODUCTION

With the development of urbanization and industrialization, the contamination of heavy metals which are directly produced and released by human activities directly implicated in the generation of oxidative stress in the plant surrounding environment which make ecological environment became unbalanced and deteriorated. Cadmium (Cd) is one of the major industrial pollutants because its highly toxicity and migration which shows its phytotoxicity even at low doses (Jabeen et al., 2009; Wahid et al., 2010).

This metal is especially active in the soil so it can be absorbed and accumulated by plant which may cause obviously damage to plant growth and development. When Cd is over accumulated by plant, it will influence plant physiology and biochemistry activity such as photosynthesis, respiration, transpiration or even relative gene expression then performed by plant appearance such as leaves chlorosis, growth inhibition and even plant death. In addition, recently reports showed that after adding nitrogen to plant, the damage caused by Cd stress can be efficiently alleviated. Moreover, plant even present better growth than those of plants which did not under stress. Some research indicate that adding appropriate nitrogen in nutrient solution can obviously enhance the development of root system and increase accumulation of Cd in root region (Liao et al., 2006; Zeng et al., 2009; Mukhopadhyay and Maiti, 2010). Further, under the same concentration of Cd stress, the biomass of *Kandelia candel* dramatically increased with the increasing nitrogen. Refer to these different research, researchers agreed that adding nitrogen can evidently alleviate the damage from Cd stress and this effect can be differential along with the different plant species as well as the different concentration of both Cd and nitrogen (Namdjoyan et al., 2011). *Nicotiana tabaccum* is famous for fast growth, ornamental value and high adaptability which is widely used as afforestation and timber tree species. However, with the development of industrialization in southwestern Tunisia, the soil suffers increasing heavy metals pollution which had detrimental influence upon the growth and development of *N. tabaccum* in this region. In this study, we comprehensively investigated the morphological, physiological and biochemical responses of *N. tabaccum* under the interaction of different concentrations of nitrogen and Cd treatment. The aims were to test the detoxification effect of nitrogen to Cd stress in *N. tabaccum* as well as to study the interaction effect of heavy metal and nutrient element on *N. tabaccum*. Therefore, the differential influence of different concentrations of Cd and N treatments on *N. tabaccum* including the growth of shoot and root, chlorophyll content, antioxidant enzymes activities and lipid peroxidation, free proline content were observed after 7 days exposure to different concentrations of N and Cd treatments. Hoping the results of this study can provide us theoretical and practical experiences to the protection of *N. tabaccum* as well as other naive plants suffered from heavy metals pollution, thus promoting the development of methodology of plant remediation (Panwan et al., 2010).

MATERIAL AND METHODS

Plant material and treatments: seeds of tobacco (*Nicotiana tabaccum*) were sterilized in 10 % (v/v) hydrogen peroxide, after imbibition, the seeds was germinated on moistened filter paper. After 7 days, uniform seedlings were transferred to plastic beakers filled with continuously aerated, basal nutrient solutions. At the age of 10 days after transplant, cadmium was added to the medium as CdCl₂ at 0, 5 and 50 µM and the 3 treatments of nitrogen comprising control (0 mM), low N (0.1 mM) and high N (5 mM). After one week of CdCl₂ treatment, plants were separated into shoots and roots, samples were stored in liquid nitrogen for subsequent analysis or dried at 70°C for at least three days in order to determine both dry material and ionic contents.

So, there were totally 9 treatments and for each treatment it was set up in triplicate to ensure the reproducibility of results. Shoots and roots samples were cut with a scissor, harvested and stored immediately at -80°C for later analysis.

Chlorophyll Concentration: chlorophyll were extracted and estimated following Arnon (1949). Absorbance of the solution was then measured at 645 and 663 nm in order to determine the concentrations of chlorophyll *a*, and chlorophyll *b*, respectively. Chlorophyll concentrations were expressed as mg g⁻¹ fresh weight.

Measurements and assays of antioxidant enzymes activities: For SOD, CAT and POD extraction, leaf samples (0.5 g fresh leaves) were ground in liquid nitrogen and extracted with 50 mM potassium phosphate buffer (pH 7.8) which contained 0.5 mM EDTA with pre-chilled pestle and mortar. Each homogenate was transferred to centrifuge tubes and was centrifuged at 15000 g, 4°C for 15 min.

The supernatant was used for enzyme activity assay. For APX extraction, leaf samples (0.5 g) were homogenized in 50 mM ice cold potassium phosphate buffer (pH 7.5) which contained 0.5 mM EDTA, 2 mM Ascorbate (AsA) and 5% Poly Vinyl Pyrrolidin (PVP) with pre-chilled pestle and mortar. Other stages were similar to extraction of other enzymes (Esfandiari et al., 2007). All operations were performed at 0-4°C. SOD activity was estimated according to Sairam *et al.* (2002), CAT activity was measured according to Aebi (1984) and the activity of POD were assayed according to the method of Guo *et al.* (2004) by monitoring the rate of guaiacol oxidation at 470 nm. In addition, APX activity was measured according to Yoshimura *et al.* (2000) by monitoring the rate of ascorbate oxidation at 290 nm. A unit of antioxidant enzymes activities was expressed as the change in absorbance per minute and specific activities as enzyme units per g of Fresh Weight (FW).

Measurement of lipid peroxidation: The level of lipid peroxidation was expressed by Malondialdehyde (MDA) concentration which was measured by colorimetric method according to Stewart and Bewley (1980). The 0.5 g of leaf samples were homogenized in 6 mL of 50 mM potassium phosphate buffer (pH 7.8). To each 1 mL aliquot of the supernatant, 2 mL of 0.6% Thiobarbituric Acid (TBA) in 10% TCA was added. The mixtures were incubated at 95°C for 15 min then quickly cooled the reaction tubes with an ice bath. After that the mixtures were centrifuged at 10,000 g for 15 min. The absorbance of the supernatant was determined at 600, 532 and 450 nm by spectrophotometer. Lipid peroxidation was expressed as the MDA content in nM per g of Fresh Weight (FW).

Measurement of free proline: Free proline was extracted and its concentration was measured by using the method of Bates *et al.* (1973). Leaf samples (0.5 g) were homogenized with 5 mL 3% sulfosalicylic acid and then the homogenate was centrifuged at 3000 g for 10 min. After that the supernatant was heated (80°C) for 1 h and then the absorbance of the supernatant was determined at 520 nm.

Statistical analysis: All data presented were means± standard value and the measurements were done with three replicates for statistical validity. One-way Analysis of Variance (ANOVA) was performed to check the variability of data and validity of the results. The results were considered significant if p<0.05.

RESULTS

Growth: The growth of shoot and root of *N. tabaccum* for 7 days exposure to treatments were shown in Fig. 1. It showed that under the same N treatment, different concentrations of Cd stress on *N. tabaccum* resulted in a significant inhibition on the growth of both shoot and root and this inhibition expressed more significantly with the increasing Cd concentrations. In addition, under the same Cd treatment, the shoot and root growth shows a dramatically increase along with the concentration of N increased. Results of univariate analysis (Table 01) indicated that Cd and N treatment both have extremely significant influence on the shoot and root growth of *N. tabaccum* ($p < 0.01$) and the interaction effect of Cd and N is significant ($p < 0.05$).

Table 1. Analysis of variance significance for the interaction effects of Cd and N on shoot and root growth in *N. tabaccum*.

Univariate	Shoot		Root	
	F	P	F	P
Cd	72.834	0.0007<0.01	31.273	0.0036<0.01
N	97.375	0.0004<0.01	19.483	0.0087<0.01

Table 2. Analysis of variance significance for the interaction effects of Cd and N on chlorophyll content in *N. Tabaccum*.

Univariate	Chlorophyll content	
	F	P
Cd	334.090	0.0001<0.01
N	87.756	0.0005<0.01

Table 3. Analysis of variance significance for the interaction effects of Cd and N on antioxidant enzymes activities in *N. Tabaccum*.

Univariate	SOD		POD		CAT		APX	
	F	P	F	P	F	P	F	P
Cd	15.162	0.0136<0.05	48.581	0.0016<0.05	27.129	0.0047<0.05	13.393	0.0169<0.05
N	11.407	0.0429<0.05	28.399	0.0043<0.05	2.777	0.1753<0.05	16.233	0.012<0.05

Table 4. Analysis of variance significance for the interaction effects of Cd and N on MDA and free proline content in *N. Tabaccum*.

Univariate	MDA		Free proline	
	F	P	F	P
Cd	69.768	0.0008<0.01	95.642	0.0004<0.01
N	61.172	0.001<0.01	3.844	0.1171>0.05

The concentrations of chlorophyll: Chlorophyll is the main pigment which can help plant to photosynthesize. When plant exposure to stress, their photosynthesis will be inhibited and the concentrations of chlorophyll can directly indicate the extent of stress-induced damage in plant. The concentrations of chlorophyll in the leaves of *N. tabaccum* for 7 days exposure were shown in Fig. 2. All Cd treatments indicated a significant decrease in chlorophyll contents. It is obvious that under the same N level, the concentrations of chlorophyll in the leaves treated with Cd significantly decreased compared with that of control which indicated that Cd treatment had significant effect on the synthesis of chlorophyll and the inhibition effect was more significant along with the increased Cd level. On the other hand, the N treatment represented a significant improvement on the chlorophyll contents. The concentrations of chlorophyll witnessed a gradual rise along with the increased N level under the same concentration of Cd stress. According to the multiple comparison (Table 02), Cd and N treatment both had extremely significant influence on the chlorophyll content of *N. tabaccum* ($p < 0.01$) and the interaction effect of Cd and N was significant ($p < 0.05$).




Fig. 1. Interactive effects of Cd and N treatment on growth of height and DBH in *N. tabaccum*. (A) Shoot growth, (B) Root growth.

The activities of antioxidant enzymes: Enzymatic antioxidant system is one of the protective mechanisms including SOD, POD, CAT and APX which can eliminate Active Oxygen Species (AOS) in plant cell and alleviate stress-induced oxidative damage (Hatata and Abdel Al, 2008). The activities of 4 antioxidant enzymes in the leaves of *N. tabaccum* for 7 days exposure were shown in Fig. 3. According to Fig. 3A, it is shown that under the same concentration of N treatment, Cd stress induced a significant increase in SOD compared with control which indicated that *N. tabaccum* has a certain degree of resistance to Cd stress.

Even when the Cd stress level reached at 5 μM , the stress didn't exceed the maximum resistance of plants. Additional, SOD activities represented a significant increase with the increasing N concentration under the same Cd level. From the results of multiple comparison (Table 03), it is indicated that Cd and N treatment both had significant influence on SOD activities ($p < 0.05$).

POD activities were significant influenced by the interaction of Cd and N treatment. According to the Fig. 3B, it is shown that under the same N level the POD activities gradually increased compared to that of control. On the contrary, at the same Cd treatment, POD activities witnessed a significant decrease along with the increased N level. From the results of multiple comparison (Table 03), both Cd and N treatment had significant influence on POD activities ($p < 0.05$).




Fig. 2. Interactive effects of Cd and N treatment on chlorophyll contents in *N. Tabaccum*.

For the CAT activities, there was a significant difference induced by different Cd treatment under the same N level ($p < 0.05$). The activities witnessed a significant growth followed by a decline (Fig. 3C) which indicated that with the increasing concentration of Cd stress, plant generated more CAT to eliminate active oxygen.

However, when the Cd stress reached to a certain degree, CAT activities were inhibited. Additional, there was no significant difference among three N concentrations under the same Cd level (Table 03). Both Cd and N treatment have significant influence on APX activities ($p < 0.05$). According to Fig. 3D, the APX activities showed a gradual rise with the increasing concentrations of Cd treatments under the same N level which indicated that Cd treatment had significant effect on the APX activities. On the contrary at the same Cd treatment, APX activities witnessed a slight decrease along with the increased N level.

The concentrations of MDA and free proline: Malondialdehyde (MDA) is the main product of membrane lipid peroxidation which can indicate the extent of stress-induced damage in plant. The results in Fig. 4A showed that the MDA concentration in the leaves of *N. tabaccum* is extremely increased under the Cd treatment compared with that of control which suggested that the Cd treatments had a significant positive correlation to the MDA accumulation ($p < 0.01$) (Table 04). On the contrary, under the same Cd level, the MDA content witnessed a significant decrease with the increasing concentration of N and the difference between different N levels was significant. Free proline plays an important role in osmoregulation and its accumulation was considered significant for the adaptation to stress in plant (Sharma and Dietz, 2006). The free proline concentrations in leaves of *N. tabaccum* showed a significant difference under




Fig. 3. Interactive effects of Cd and N treatment on antioxidant enzymes activities of *N. tabaccum*, (A) SOD activities, (B) POD activities, (C) CAT activities, (D) APX activities.

Cd stress ($p < 0.01$). With the increasing Cd level, free proline content had a rapid rise followed by a slowly increase which indicated that Cd stress improved the accumulation of free proline and played a protective role in *N. tabaccum* (Fig. 4B). However, when the Cd level reached to a certain degree, the synthesis of proline was inhibited and influenced the increase of proline. Additional, N treatment had no significant influenced on the concentration of free proline under the same Cd level and there was also no significant difference among three N concentrations, which indicated that the concentrations of free proline were only affected by Cd stress.

DISCUSSION

Cd contaminated has detrimental influence on plant growth and development. Over-accumulation of Cd concentration in plants shows differential symptoms of toxicity including the growth inhibition, leaves chlorosis and biomass decrease. The results showed that Cd stress obviously inhibited the growth of *N. tabaccum* slowed down the synthesis process of chlorophyll which decreased its concentration in plants.




Fig. 4. Interactive effect of Cd and N treatment on MDA and free proline content in *N. tabaccum*, (A) MDA content, (B) Free proline content.

Over-accumulated Cd injury to plants is mainly because of altered oxidant levels which causing the occurrence of oxidative stress due to accumulation of Active Oxygen Species (AOS) including superoxide radical (O_2^-), Hydroxyl radical (OH^-) and hydrogen peroxide (H_2O_2). In normal circumstances, the speed of generation and elimination of AOS is balanced. However, when plants exposed to Cd stress, plants generate more AOS than elimination which cause over accumulation of AOS in plants which lead to significant damage to membrane system in plants and induce lipid peroxidation, increase permeability and conductivity (Smiri et al., 2010; Xu et al., 2011, Nayari et al., 2012). The concentration of Malondialdehyde (MDA) which is a general indicator of lipid peroxidation can demonstrate the extent of oxidative stress in plants (Xu et al., 2011). On the contrary, plant cells have their own defense mechanisms against AOS. One of the protective mechanisms is the enzymatic antioxidant system which involves the sequential and simultaneous action of a number of enzymes such as SOD, POD, CAT, APX (Gill et al., 2011; Mao et al., 2011). The results showed that under Cd stress, MDA and free proline concentration in plants was significant increased with the increasing Cd level. However, *N. tabaccum* showed a certain degree of resistance to Cd stress and the concentration of Cd treatment didn't exceed its maximum capacity of resistance. Thus the oxidative stress caused by Cd addition was moderate which allowing the plants to adapt it actively by increasing their own antioxidant enzymes activity. Meanwhile, elevated level of these antioxidative enzymes in plants following the Cd treatment in this study demonstrated that these enzymes acted together to alleviate the impact of Cd stress. Furthermore, it is worth noting that the influenced of Cd stress on plant has significant difference among those three Cd concentrations.

Nitrogen is one of the essential nutrient elements. The deficiency of nitrogen will lead to metabolic disturbance in plants thus causing the decrease of plant growth and productivities. With pollution of heavy metals being serious day by day, the heavy metals stress may become the main reason of the deficiency or inefficiency of nutrient element. Meanwhile, supplement of nitrogen for plants can obviously alleviate heavy metal-reduced damage. Cela and Sumner (2002) suggested that adding nitrogen to soil, nitrification will occur and obviously decreased the pH of soil thus significant increasing the solubility of heavy metals in soil which decrease the adsorption amount of heavy metals.

CONCLUSION

In this study, the researchers discovered that nitrogen could effectively alleviated Cd induced damage in *N. tabaccum* and there was a significant difference of effects on plants between lower N concentration (0.1 mM) and higher N concentration (5 mM). Supplement of high-concentration nitrogen to *N. tabaccum* under Cd stress obviously enhanced plants growth, promoted the synthesis of chlorophyll as well as the activities of antioxidant enzymes which partially alleviated the accumulation of AOS associated with Cd exposure which indicated that the increasing concentration of N supplement had more significant effect of alleviation to Cd induced damage in *N. tabaccum*.

REFERENCES

- Abei H. 1984. Catalase in vitro. *Methods Enzymol* 105: 121-126.
- Arnon DI. 1949. Copper enzyme is isolated chloroplast polyphenol oxidase in Beta vulgaris. *Plant Physiol* 24: 1-15.
- Bates, L. S., R. P. Waldren, and I. D. Teare. 1973. Rapid determination of free proline for water-stress studies. *Plant Soil* 39: 205-207.
- Cela, S. and M. E. Sumner. 2002. Soil Zinc Fractions Determine Inhibition of Nitrification. *Water, Air & Soil Pollution* 141(1-4): 91-104.
- Esfandiari EO, Shakiba MR, Mahboob SA, Alyari H, Toorchi M, Water stress, antioxidant enzyme activity and lipid peroxidation in wheat seedling. *Int. J. Food Agric. Environ* 5: 149-153. 2007.
- Gill SS, Khan NA, Tuteja N, Differential cadmium stress tolerance in five Indian mustard (*Brassica juncea* L) cultivars. An evaluation of the role of antioxidant machinery. *Plant Signal Behav* 6(2): 293-300. 2011.
- Guo T, Zhang G, Zhou M, Wu F, Chen J, Effects of aluminum and cadmium toxicity on growth and antioxidant enzyme activities of two barley genotypes with different Al resistance. *Plant Soil* 258: 241-248. 2004.
- Hatata MM, Abdel Aal EA, Oxodative stress and antioxidant defense mechanisms in response to cadmium treatments. *American-Eurasian J. Agri. Env. Sci* 4(6): 655-669. 2008.
- Jabeen R, Ahmad A, Iqbal M, Phytoremediation of heavy metals: Physiological and Molecular Mechanisms. *Bot. Rev.* 75(4): 339-364. 2009.
- Liao H, Wan H, Shaff J, Wang X, Yan X, Kochian LV, Phosphorus and aluminum interactions in soybean in relation to aluminum tolerance. Exudation of specific organic acids from different regions of the intact root system. *Plant Phys.* 141: 674-684. 2006.
- Mao S, Robiul MI, Hu Y, Qian X, Chen F, Xue X, Antioxidant enzyme activities and lipid peroxidation in corn following soil application of superabsorbent polymer at different fertilizer regimes. *African J. Biot* 10(49): 10000-10008. 2011.
- Mukhopadhyay S, Maiti S.K, Phytoremediation of metal mine waste. *App. Eco. Env. Resea* 8(3): 207-222. 2010.
- Namdjoyan SH, Khavari-Nejad RA, Bernard F, Nejdassattari, Shaker T, Antioxidant defense mechanisms in response to cadmium treatments in two sanflower cultivars. *Russian J. Plant Phy* 58(3): 467-477. 2011.
- Nazari M, Maali Amiri R, Mehraban FH, Khaneghah HZ, Change in Antioxidant Responses against Oxidative Damage in Black Chickpea Following Cold Acclimation. *Indian J. Bot* 59(2): 209-215. 2012.
- Panwar BS, Marton L, Kdar I, Anton A, Nemeth T, Phytoremediation, a novel green technology to restore soil health. *Acta Agronomica Hungarica* 58(4): 443-458. 2010.

- Rastgoo L, Alemzadeh A, Biochemical response of Gouan to heavy metals stress. *Australian J. Crop. Sci* 5(6): 457-470. 2011.
- Sairam RK, Roa KV, Srivastava GC, Differential response of wheat genotypes to long term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration. *Plant Sci* 163: 1037-1046. 2002.
- Sharma SS, Dietz KJ, The significance of amino acids and amino acid-derived molecules in plant responses and adaptation to heavy metal stress. *J. Exp. Bot* 57(4): 711-726. 2006.
- Smiri M, Chaoui A, Ferjani EE, Interaction Between Heavy Metals and Thiol-linked Redox Reactions in Germination. *J. Biol. Sci* 13: 877-883. 2010.
- Stewart RRC, Bewley JD, Lipid peroxidation associated aging of soybean axes. *Plant Physiol* 65: 245-248. 1980.
- Wahid A, Arshad M, Farooq M, Cadmium Phytotoxicity: Responses, Mechanisms and Mitigation Strategies. *Sustainable Agri. Rev.* 1: 371-403. 2010.
- Xu HW, Lu Y, Tong SY, Song FB, Lipid peroxidation antioxidant enzyme activity and osmotic adjustment changes in husk leaves of maize in black soils region of Northeast China. *African J. Agr. Resea.* 6(13): 3098-3102. 2011.
- Yoshimura K, Yabute Y, Ishikawa T, Shigeoka S, Expression of spinach ascorbate peroxidase isoenzymes in response to oxidative stresses. *Plant Physiol* 123: 223-233. 2000.
- Zeng X, Ma LQ, Qui R, Tang Y, Responses of non-protein thiols to Cd exposure in Cd hyperaccumulator *Arabis paniculata* franch. *Env. Exp. Bot* 66: 242-248. 2009.

Corresponding author: Dr. Chiraz Chaffei Haouari, Université Tunis El Manar, Faculté des Sciences de Tunis, Département de Biologie, 1060-Tunis-TUNISIE Fax: 216 71 885 480. Email: chiraz_ch2001@yahoo.fr