Irrigation Methods, Physiological and Phenological Characteristics of Pomegranates and Pomegranate Trees

^{Ву} Hamid Kheyrodin and Khosro Ghazvinian

ISSN 0970-4973 (Print) ISSN 2319-3077 (Online/Electronic)

Volume 30

No.1 (2013)

J. Biol. Chem. Research Volume 30 (1) 2013 Pages No. 27-44

Journal of Biological and Chemical Research

(An International Journal of Life Sciences and Chemistry)

Published by Society for Advancement of Sciences®

J. Biol. Chem. Research. Vol. 30, No. 1: 27-44 (2013) (An International Journal of Life Sciences and Chemistry) Ms 30/1/28/2013, All rights reserved <u>ISSN 0970-4973 (Print)</u> <u>ISSN 2319-3077 (Online/Electronic)</u> Published by Society for Advancement of Science[®]



JBCR

http:// <u>www.jbcr.in</u> jbiolchemres@gmail.com <u>info@jbcr.in</u>

RESEARCH PAPER Received: 06/11/2012 Revised: 02/01/2013 Accepted: 04/01/2013 Irrigation Methods, Physiological and Phenological Characteristics of Pomegranates and Pomegranate Trees *Hamid Kheyrodin and **Khosro Ghazvinian

*Faculty of Desert Science, Semnan University, Iran **Faculty Animal Science, Semnan University, Iran

ABSTRACT

The main irrigated garden Pomegranate pomegranates are in quantitative yield. Note that the climate is generally warm and dry pomegranates are grown. The level of annual evaporation is about 2500 mm and the boundary of the water as hot and dry trees also act as a substantial amount of transpiration and evaporation of water in the soil is lost. This is the fruit of the pomegranate, which requires a long growing season and irrigation on tree increases its sensitivity. In this case, there are different numbers that are not generally relevant experience and product evaluation of the theoretical and academic. We discussed what we needed at least one tree produces around 25 tons per hectare and economic exploitation is dependent on experience. It is dry at low water, and Iran can only save water, and reliance on the optimal irrigation efficiency is not high. Daily water requirement in the following table pomegranate tree in a garden with a density of 1,000 trees per hectare and canopy trees in the garden are given by 50 percent of the irrigated area. Many fruit trees are not very sensitive to soil texture.

If soil depth and drainage are adequate and proper acidity might have grown up to be good enough to make the product. It is necessary to note that soils are usually lighter than the heavy soils with low humidity and warm earlier in the spring so cold later in the fall. And it also has a better ventilation if adequate water and nutrients to grow more fruit trees.

Key words: Pomegranate, remote sensing, irrigation systems performance.

INTRODUCTION

Benefit to Trees

The increased use of underground irrigation systems, including the growing use of the drip-irrigation method, has greatly expanded the variety of plant materials that can be used in the community landscape. In addition to turfgrass, shrubs and trees can benefit from the use of a well-designed and operated irrigation system. Irrigation of trees should be a component of a good Plant Health Care plan and should be considered when developing a strategy for the care and maintenance of trees growing in any landscape. Since drought stresses are often cyclical in many regions of the country, their occurrence should be planned for during the development of any tree management strategy.

Lack of moisture is one of the more severe problems that affect trees growing in a landscape. Trees growing in urban conditions are often more susceptible to this damage, due to reduced soil volumes, poorer soil structure and desiccation. However, trees in any landscape need adequate moisture to survive, and underground irrigation systems can enhance the opportunity for survival. Irrigation of trees can provide benefits that lead to healthier trees, which are more able to withstand other stresses such as insect and disease infestations that might occur. Reducing drought stress through the proper use of an irrigation system can greatly enhance the likelihood of newly planted trees reaching maturity, and can provide older trees with a consistent source of moisture, even during droughty periods.

Irrigation water to crops is an effective means of enhancing agricultural production and productivity. In addition, irrigation provides the basis for a better and more diversified cropping pattern and growing of high-value crops, and thus facilitates overall improvement in socio-economic conditions of the farming community. A considerable amount of work has been undertaken in the past 15 years to develop a framework for irrigation performance assessment. Irrigation performance indicators cover the traditional aspects related to adequacy, equity and reliability of the water service (Wolters 1992; Murray-Rust & Snellen 1993; Bos et al. 1994). The framework of performance indicators includes environmental sustainability features to evaluate the longer-term effects irrigation has on the environment such as groundwater table changes (Bos et al. 1991). The literature reported in the last two decades indicates that the performance of irrigation schemes, particularly large areas, is far below expectations. The magnitude of underperformance is well represented by Seckler et al. (1988). Up to the mid-1980s, the lack of performance indicators was the main reason preventing the evaluation of irrigation system performance in guantifiable units. Moreover, monitoring and evaluation have received much lip service in the present decade and have seldom been carried out properly and effectively. One of the main reasons is that the conventional methods of data collection through field observation are difficult, timeconsuming and cannot be carried out at the same time particularly in large irrigated commands. In addition, monitoring is not possible at frequent intervals. ISP assessment based on conventional data. The evaluation of the efficiency of irrigation water use has undergone major modifications during the last 25 years.

It moved from the classical irrigation efficiencies (Bos & Nugteren 1974; Jensen 1977) to the current performance indicators (Levine 1982; Small & Svendsen 1990; Bos et al. 1994) and, to the framework of water accounting at a regional scale (Molden 1997; Burt et al. 1997). The required data to calculate all these indicators consist of discharge measurements, crop irrigation water requirements, effective rainfall, actual evapo-transpiration, irrigated area, cropping intensity and crop yield among others.

ISP assessment based on remote sensing data. Remote sensing data to determine actual evapotranspiration and crop water stress for managing irrigation systems was suggested by several authors in the eighties (e.g. Jackson et al. 1981; Jackson 1984; Ben Asher et al. 1986). However, Azzali & Menenti (1987) showed how high resolution Landsat images could be used to map crops and crop phenology stages in two Italian irrigation districts, and to detect crop-specific effects such as water stress. Menenti et al. (1989) made a list of indicators that could be quantified with remotely sensed data. ISP assessment based on physiological data. The use of physiological traits as screening tools in breeding for yield is still largely experimental, for different reasons. Sometimes the traits are very indirectly related to yield or there is little ecophysiological understanding of the crop, especially when breeding forvield under stress. Nevertheless, breeding for crop escape has been very successful; phonological changes have been the most important indirect factor in increasing wheat yields under Mediterranean conditions. However, in breeding for crop resistance, the evaluated traits and screening tools are often related to tolerance, not avoidance. An indirect (i.e. physiology-based) breeding strategy could fail to produce yield gains and might even lead to a decrease in yield. Improved plant tolerance, though it protects the crop, can limit yield potential. The most promising ecophysiological methods allow for quick screening of "integrative" physiological traits, so called because they integrate physiological processes either in time (i.e. during the plant cycle) or at the organization level (e.g. whole plant, canopy). Nevertheless, remote sensing detection of fluorescence spectra at the canopy level could become a promising approach for breeding purposes. Despite the objections shown for breeding strategies for yield, the physiological tools will ultimately be related to the performance of the irrigated crop. We propose carbon isotope composition of plant tissues and spectral reflectance methods, as integrative physiological traits that could be used in assessing the performance and the hydrological efficiency of irrigation systems, in order to improve water saving.

PHYSIOLOGICAL TOOLS

The physiological tools presented here were originally proposed to small cereal grains breeders interested in adopting a physiological approach to the improvement of their varieties. The potential contribution of the physiological approach to plant selection, as well as their inherent limitations and the requirements to fit, have been recently reviewed extensively from a purely breeding perspective (for example, Jackson et al. 1996). The theoretical frameworks to define the physiological determinants of yield of cereal crops which are the obvious candidates to be evaluated have also been reasonably well established.

One approach to search for 'potential' traits to be used in breeding programs is to identify the physiological processes determining productivity. Therefore, yield can be easily divided in several integrative components or traits. The yield itself is the most integrative trait, because it is influenced by all the (known and unknown) factors which are actually determining productivity. However, there many known limitations in a purely empirical approach to breeding only based in yield. Therefore, any breeding strategy based in a physiological (i.e. analytical) approach should use screening tools or criteria that would allow evaluation the integrative physiological parameters which are determining harvestable yield with a single measurement. For the assessment of the performance of the Irrigation Systems, it should be defined a set of suitable physiological criteria of the crops, easy to be measured with the proposed tools, and with strong correlation with the classical irrigation performance indicators.

Damage to Trees

While underground irrigation systems can provide a useful benefit to establishing and growing trees in the landscape, often the installation and improper use of these systems leads to serious tree health issues. Root damage and overwatering are the most common problems associated with underground systems that are installed to irrigate trees or turf grass.

Root Damage

The long term health of trees can be affected during by the installation of underground irrigation systems as a result of trenching near a tree. Research has shown that tree roots can travel a distance away from the trunk, over 2 1/2 times the height of the tree, and that over 85% of the root growth occurs within 18 inches of the soil surface. Often these roots are severely damaged, or cut entirely, during the installation of the underground piping that makes up the irrigation system. In many cases trenching, vibratory plowing or earth sawing operations make vertical cuts through the soil to depths that can range to over 20 inches, cutting many tree roots growing in their path. The damage to trees is not readily apparent, but generally begins to appear within months or years of the damage. Dieback, decline and mortality often occur, but the severing of roots during the irrigation installation process is not recalled, since most of the problem took place underground, out of sight of the property owner or contractor. Arborists can provide advice to irrigation installation contractors on how to minimize damage to trees during the installation process. Arborists should be used in developing a tree protection plan that will minimize potential impacts to the root zones of the trees during the installation of any underground system. Without planning for the protection of trees growing in the vicinity of the installation of an underground irrigation system, tree root damage is likely to occur. The layout, installation and operational plan for the underground irrigation system should be reviewed by an arborist. These reviews provide the opportunity to protect trees and ensure that their healthy growth will continue. One of the most useful, and tree friendly, techniques for installing irrigation or other utility piping in the vicinity of trees is horizontal directional drilling.

This "trenchless technology" involves the use of specialized equipment which uses a boring head to tunnel under the root zone of a tree, limiting any potential impact to the tree's underground life support system. Since most tree root growth occurs in the top 18 inches of soil, directional drilling is used to pull piping or conduit at depths, which in most cases, are safely below the root zone.

There are several sources of directional drilling equipment and a variety of models available, depending on the specific application. Units range in size from self contained, small footprint sized units, to larger systems that are used for longdistance utility and telecommunication installations. Today, affordable trenchless technology tools are available from a variety of manufacturers and wide ranges of product specific features are available. Small, easy to maneuver units are extremely versatile and can be used for a variety of applications related to irrigation.



Overwatering

Another serious problem affecting trees as a result of underground irrigation systems is overwatering and soil saturation. Many times an irrigation system is installed and then programmed for "automatic" operation. This often results in overwatering the root zone of trees, resulting in reduced oxygen saturation or shallow root growth. Since many irrigation systems are designed and installed primarily to water turfgrass, the effect on nearby trees is not considered.

This often results in watering practices that are detrimental to healthy tree root growth. Additionally, overwatering can result in shallow root growth, resulting in less healthy trees that are more prone to a variety of problems including wind throw and reliance on artificial watering regimes. However, it is possible for trees and turfgrass irrigation regimes to coexist successfully. Proper watering for good turf growth can compliment healthy tree root growth. Deeper and less frequent watering can encourage healthier root growth for both turf and trees. Today, there are a variety of opportunities for arborists to become a more integral component in the design of underground irrigation systems. In today's marketplace, tree professionals can provide valuable professional expertise, knowledgeable installation services and a new perspective to underground irrigation construction operations. Helping to protect trees ensures healthy growth and limits the potential impacts due to underground activities. This is a key role that is best filled by a Massachusetts Certified Arborist.

RESULTS AND DISCUSSION

Another example of the various species and compounds where $\Delta 13C$ analyses are applicable is shown in Fig. 2b. Ferrio et al. (2003a) studied the $\Delta 13C$ from wood samples of Aleppo pine (Pinus halepensis) and holm oak (Quercus ilex). Both species are typically Mediterranean, but they display contrasting strategies against drought. The former is a clear drought avoiding species, with a fast stomatal closure, whereas the latter relies mostly on a very effective water uptake, and is partly drought tolerant. They also found a strong relationship between $\Delta 13C$ and annual precipitation (unlike grains, tree rings are formed throughout the year), which was also steeper among the most arid environments. Interestingly, $\Delta 13C$ in Aleppo pine was far more sensitive to water availability than in Holm oak. Again, the explanation relies on the close relationship between $\Delta 13C$ and stomatal conductance. As the stomata of Aleppo pine showed faster responses to drought, the $\Delta 13C$ for this species have greater variations with changes in water availability. In contrast, the Holm oak can extract water from drier soils, as well as from deeper water tables, showing relatively low variations in $\Delta 13C$ among trees growing in contrasting environments. Thus, although the relationship between $\Delta 13C$. The pattern of light reflection on leaves at different wavelengths through the photosynthetically active radiation (PAR, 400-700 nm) and near infrared radiation (NIR, 700-1200 nm) regions of the electromagnetic spectrum is very different from that of soil and other materials (Figure 3).

Leaf pigments absorb light strongly in the PAR region but not in the NIR, thus reducing the reflection of PAR but not of NIR. Such a pattern of pigment absorption determines the characteristic reflectance signature of leaves (Figure 4). Similarly, the light spectrum reflected by a canopy (either natural or agricultural) differs from that reflected by the bare soil and varies in a way that can be related to the overall area of leaves and other photosynthetic organs in the canopy, as well as to their pigment composition and other physiological factors (Figure 5).



Figure. 1. Simplified scheme of the relationship between carbon isotope discrimination (Δ 13C) in C3 plants and stomatal conductance. A) High stomatal conductance, high discrimination: CO2 diffusses easily into the intercelular space, the activity of the carboxylating enzyme (represented as R) is not limited by CO2 concentration and thus it has more chances to discriminate against 13C. B) Low stomatal conductance, low discrimination: the flux of CO2 is reduced, and the limiting factor of photosynthesis is stomatal conductance. In this case RuBisCO is forced to fix a higher proportion of 13C. Redrawn from Ferrio et al. (2003b)



Figure. 2. Relationship between: a) water inputs (WI, rainfall plus irrigation if applied) during grain filling and Δ 13C of barley and wheat grains. Data from Araus et al. (1997a) for barley and from Araus et al. (1999; 2003) for wheat; Δ 13Cbarley= 9.99 + 1.52*In (WI); Δ 13Cwheat= 8.50 + 1.78*In (WI) b) mean annual precipitation of 20-25 years and Δ 13C of wood from the corresponding tree-rings of Aleppo pine and holm oak. Redrawn from Ferrio et al. (2003a); Δ 13Cpine= 4.63 + 1.91*In (WI); Δ 13Coak= 11.72 + 0.97*In (WI).

Therefore, the measurement of spectra reflected by vegetation canopies provides information that can be used to estimate a large scope of parameters. Some of them are related to the green biomass of the canopy, its photosynthetic size (i.e. total area of leaves and other photosynthetic organs), the amount of PAR absorbed by the canopy, and its photosynthetic potential. Other parameters are more related to the canopy's physiological status at the time of measurement and can be used to assess the extent of some nutrient deficiencies and environmental stresses. The physiological parameters that can be estimated by spectral reflectance techniques include chlorophyll and carotenoid concentrations, photosynthetic radiation use efficiency (PRUE), and water content. Water Index (WI) has been used to track changes in a number of parameters related with the water status of the crop (leaf water potential, stomatal conductance, and other). Some of these physiological parameters might show a kind of relationships with indicators of classical irrigation efficiencies used for assessment of ISP, in a similar manner than satellite measurements.

Spectral reflectance indices are formulations based on simple operations between reflectance at given wavelengths, such as ratios, differences, etc, which are widely used to quantitatively relate changes in reflectance spectra to changes in physiological variables. These indices have the advantage of summing up in a few numbers the large amount of information contained in a reflectance spectrum with narrow waveband resolution.



Figure. 3. Spectral reflectance from crop surfaces.



Figure. 4. Reflectance signature of two wheat leaves differing in nitrogen status. Note the higher reflectance in the PAR region of the nitrogen deficient leaf due to lower chlorophyll content in the leaf area.

Originally used in remote sensing by aircraft and satellites, reflectances measured at the ground level are very useful for assessing agrophysiological traits. These traits can be evaluated simultaneously in each sample, at a rate of up to one thousand samples per day, which is much more tedious and time consuming with other methods. This makes spectroradiometric indices ideal for screening for yield potential or for resistance to different stresses, and eventually for assessment of ISP. *Sample applications* Perhaps the most widespread application of reflectance indices is for assessing parametersrelated to canopy greenness.

These parameters are related to the canopy's photosynthetic size and include green biomass, leaf area index (LAI) (total one-side leaf area of the crop relative to soil area), green leaf area index (GLAI) (similar to LAI, but includes only functional green leaves), and green area index (GAI) (similar to GLAI, but includes other photosynthetic organs such as green stems). The amount of green area in a canopy determines PAR absorption by photosynthetic organs, which in turn determines the canopy's potential production. The fraction of the incident PAR that is absorbed by the canopy (fPAR) can be estimated from LAI-related parameters or directly from reflectance measurements. Cumulative PAR absorption, which is one of the parameters determining total biomass and thus final yield, can be assessed by measuring reflectance periodically during the growth cycle. Some physiological parameters can also be quantified by spectral indices. Leaf pigments can be detected and quantified based on reflectance spectra and can be used as indicators of several physiological processes. Thus, the canopy's nutritional state can be evaluated through pigment concentration, as chlorophyll (Chl) concentration in leaves is (usually) closely correlated to its nitrogen content. Indices that are good indicators of ChI are (usually) also good indicators of N-content. In addition, plants with low N usually have a high carotenoid (Car) to Chl ratio, which can also be assessed by reflectance indices (Figure 5).



Figure. 5. Changes in the pattern of canopy reflectance of a durum wheat grown in the field.

Measurements were taken every three days (a, b, c), during the last week of the grain filling period, coinciding with fast crop senescence. Pigment remote sensing can also be used for assessing the crop's phenological stage (Figure 5) and the occurrence of several stress factors (Blackburn 1998; Peñuelas 1998). For example, the Car to Chl ratio can be associated with senescing processes that result from the plant's natural ontogeny pattern or are triggered by different stresses. Also, phenological stages can be associated with different Car/Chl values. Several indices related to changes in pigment composition have beendeveloped and can be used for the remote detection of nutrient deficiencies, environmental stresses, pest attacks, etc.

In such contexts, by periodically assessing leaf area, leaf area duration (LAD) can also be used as an indicator of resistance to certain environmental stresses.

The photosynthetic capacity of a canopy can be estimated by using vegetation indices that correlate to the photosynthetic size of the canopy or indices related to the amount of chlorophyll.

However, actual photosynthesis may not match photosynthetic capacity due to the variability of photosynthetic use efficiency of the absorbed radiation, especially when plants are exposed to unfavorable conditions (climate, performance or the irrigation system). The photochemical reflectance index (PRI) was developed to detect pigment changes in the xantophyll cycle associated with changes in PRUE (Filella et al. 1996). PRI has been shown to track the changes in PRUE induced by factors such as nutritional status and midday reduction, across different species and functional types.

Another potential application of reflectance indices is remote detection of relative water content (RWC) of plants. Different levels of water stress, which could be related with the performance of the irrigation system, can be detected indirectly through their effects on vegetation indices related to leaf area, pigment concentration, or photochemical efficiency. Specific indices like Water Index have been developed for the direct assessment of RWC.

Use of Canopy Reflectance Indices

Assessing the photosynthetic size of canopies using vegetation indices

Vegetation indices (VI) estimate parameters related to the photosynthetic size of a canopy based on the reflectances in the red and near infrared regions. Green biomass, LAI, GAI, GLAI, fPAR, etc, can be estimated through their positive correlation (either linear or logarithmic) with vegetation indices (Wiegand and Richardson 1990a, 1990b; Baret and Guyot 1991; Price and Bausch 1995). Measuring vegetation indices periodically during the crop growing cycle allows the estimation of LAD (which can be used as an indicator of environmental stress tolerance) and the total PAR absorbed by the canopy, which is one of the most important factors for predicting yield (Wiegand and Richardson 1990a, 1900b).

Vegetation indices take advantage of the great differences in reflectance at red and NIR caused by vegetation. The most widely used VI are the simple ratio (SR) and the normalized difference vegetation index (NDVI), which are defined as: SR = RNIR / RRed, with a range of 0 to ∞ , where RNIR is the reflectance at NIR and RRed is the reflectance at red. NDVI = (RNIR - RRed) / (RNIR + RRed), with a range of -1 to 1. SR and NDVI were originally used with the wide wavebands of former radiometers (for example, 550-670 nm for red and 710-980 nm for near infrared in AVHRR radiometers in satellites of NOAA series). With the high spectral resolution of today's radiometers, wavebands can be much narrower. Hall et al. (1990) used a waveband centred at 770 nm for NIR and another at 660 nm for red, while Penuelas et al. (1997b) used 900 nm and 680 nm for NIR and red, respectively. Some authors have reported improvements in NDVI performance after changing the wavebands used in the index. Carter (1998) describes an improved correlation with leaf photosynthetic capacity when using a modified NDVI where R701 (+/-2nm) and R520 (+/-2nm) were used for NIR and red, respectively.

Variations of these indices have been proposed to compensate for the effect of soil background, which might be very different depending on the crop type, and on the irrigation system. Thus the soil adjusted vegetation index (SAVI) was defined by Huete (1988) as: SAVI = [(RNIR - RRed) / RNIR + RRed + L)] (1 + L), where the parameter L was adjusted to minimize noise caused by soil for a large range of soil covers. For most crop conditions L=0.5, while for very low soil covers L=1 would be more appropriate, and L=0.25 would be appropriate for very high covers (Huete 1988). Other indices include parameters obtained from the soil's reflectance spectrum. One of them is the transformed soil adjusted vegetation index (TSAVI) which was defined by Baret and Guyot (1991) as: TSAVI = a (RNIR aRRed - b) / [RRed + a (RNIR - b) + 0.08(1+a2)], where A is the slope and b is the intercept of the linear equation RNIRsoil = a*RRed soil + b. An important drawback in estimating LAI by VI is the saturation of the VI with LAI. Saturation of NDVI starts at about LAI=1, and beyond LAI=2 it becomes insensitive to further increases in LAI (Gamon et al. 1995). Perpendicular vegetation index (PVI) partly overcomes the saturation problem inherent to NDVI (Richardson and Wiegand 1977): PVI = {(RRed.soil-RRed.vegetation) 2 + (RNIR.vegetation-RNIR.soil)21/2 Although PVI is more sensitive than NDVI to changes in the viewing geometry, PVI does not become as clearly saturated as NDVI with changes in GLAI (Shibayama et al. 1986). Examples of assessing LAI-related parameters by VI can be found in the literature (Baret and Guyot 1991; Field et al. 1994; Price and Bausch 1995). Ground level measurement of VI has been used successfully as a tool for assessing early biomass and vigour of different wheat genotypes (Elliott and Regan 1993; Bellairs et al. 1996) and for assessing the intensity of different plant stresses, Penuelas et al. (1997b). A practical use of vegetation indices is for making yield predictions. Yield can be predicted from successive VI measurements taken during the growing season, based on the following assumptions (Wiegand et al. 1991, Wiegand and Richardson 1990b, Rudorff and Batista 1990): 1) plant stands integrate the growing conditions experienced and express net assimilation achieved through the canopy, 2) stresses severe enough to affect economic yield will be detectable through their effects on crop development and the persistence of photosynthetically active tissue in the canopy, 3) high economic yields cannot be achieved unless plant canopies fully utilize available solar radiation as the plants enter the reproductive stage, and 4) vegetation indices calculated from remote observations in appropriate wavelengths effectively measure the photosynthetic size of the canopy.

Remote sensing of pigments

Estimation of chlorophyll concentration, several indices has been developed for estimating Chl concentration using canopy reflectance methods. The simplest indices are just reflectance at 675 and 550 nm. Reflectance at 675 nm (R675) is very sensitive to changes in Chl content. However, the relationship becomes saturated at relatively low Chl values (around 10 μ g cm-2) and is a good indicator of chlorophyll content only at very low concentrations. Absorption by Chl at 550 nm is lower than at 675 nm; therefore, the reflectance at this wavelength (R550) is less sensitive to changes in Chl content but is not saturated at such low concentrations, thus covering a range of higher Chl values (Thomas and Gausman 1977; Jacquemoud and Baret 1990; Lichtenthaler et al. 1996).

Both R675 and R550 are non-normalized indices that can be affected by external factors (Curran 1983). Other indices use more than one wavelength. Analyzing wavelengths that were more sensitive to changes in Chla, Chlb, and Cars in soybean leaves grown at different N levels, Chapelle et al. (1992) developed the ratio analysis of reflectance spectra (RARS) indices (RARSa = R675 / R700; RARSb = R675 / (R650 * R700); RARSc = R760 / R500), which optimized the estimation of Chla, Chlb, and Cars, respectively, in soybean leaves. Other reflectance indices that can be used for estimating pigment concentration are summarized in Table 1. In addition to the wide variety of indices related to absolute Chl concentration, the normalized phaeophytinization index (NPQI) can be used to detect chlorophyll degradation. NPQI = (R415 - R435) / (R415 + R435) (Penuelas et al. 1995c) NPQI was introduced as an indicator of pest attacks on apple trees (Penuelas et al. 1995c). Estimation of the Carotenoid to chlorophyll ratios, estimating the Car: Chl ratio by reflectance indices can be useful for assessing the extent of some plant stresses and eventually the performance of the irrigation system, given that increases in Cars concentration relative to Chl are often observed when plants are subjected to stress (Young and Britton 1990). Both Chl and Car absorb in the blue, but only Chl absorbs in the red. Indices that are combinations of the reflectance in these two regions are correlated to the Car: Chl ratio. The simplest indices are pigment simple ratio (PSR) and normalized difference pigment index (NDPI), which are formulated in an analogue way to SR and NDVI and defined to estimate the ratio of total pigments to Chla (Penuelas et al. 1993a): PSR = R430 / R680, NDPI = (R680 -R430) / (R680 + R430) Both PSR and NDPI are affected by disrupting effects introduced by leaf surface and structure. A new index was developed to avoid such problems: the structural independent pigment index (SIPI), which was defined by Penuelas et al. (1995a) as: SIPI = (R800 - R435) / (R415 + R435) SIPI uses wavelengths showing the best semiempirical estimation of the Car: Chla ratio, and its formulation minimizes the disrupting effects of leaf surface and mesophyll structure (Penuelas et al. 1995a). R800 is used as a reference where neither Cars nor Chl absorb and are only affected by the structure. Indices related to the Cars: Chl ratio change during the crop growing cycle. They are low during vegetative growth and start to increase before the beginning of senescence (Filella et al. 1995). They can be used in assessing the nutritional state of a crop (Filella et al. 1995), shown by high values of the indices when N is low, and for detecting pest attacks (Penuelas et al. 1995c).

Assessing radiation use efficiency by PRI

Canopy photosynthesis can be roughly estimated based on the estimation of the canopy's photosynthetic size or Chl concentration. However, these parameters are associated with potential canopy photosynthesis, which does not always correspond to actual photosynthesis, especially for plants growing in stressful environments. While VI are correlated with PAR absorption by the canopy (a slowly varying trait, in a range of days to weeks), the photochemical reflectance index (PRI) is correlated with photosynthetic radiation use efficiency (PRUE) of absorbed PAR, a rapidly varying process, in a range of hours (Gamon et al. 1997). Part of the PAR absorbed by Chl cannot be used for photosynthesis and is lost mainly through heat dissipation, which is linked to the xanthophyll-de-epoxidation cycle (Demmig-Adams and Adams 1996).

Irrigation.....trees

PRI reflects changes in reflectance of around 531 nm, which have been associated with pigment changes in the xanthophylls cycle (Gamon et al. 1992; Penuelas et al. 1995b). It is defined as: PRI = (R531 - R570) / (R531 + R570), where PRI is correlated with the deepoxidation stage of the xanthophylls cycle, with zeaxanthin, and with radiation-use efficiency (Filella et al. 1996). Higher PRI values indicate greater efficiency.

Directly assessing plant water status

Some bands of radiation absorption by water exist in the 1300-2500 nm region, but due to its high absorptance in this region, reflectance becomes saturated (i.e. it does not respond to further increases in RWC) even in a canopy with low water content. In the 950-970 nm regions, there is some weak absorption of radiation by water that is not saturated for a moderately dry canopy. The reflectance at 970 has been used in the definition of the water index (WI). WI = R900 / R970, (Penuelas et al. 1993b, 1997a) In WI, reflectance at 970 nm is taken as a wavelength sensitive to water content, while reflectance at 900 nm is taken as a reference which is similarly affected by canopy and leaf structures but with null absorption by water. WI has been used to track changes in RWC, leaf water potential, stomatal conductance, and foliage minus air temperature differences when plant water stress is well developed (RWC<0.85) (Penuelas et al. 1993b). Penuelas et al. (1997a) reported a correlation coefficient of around 0.55 between WI and RWC for a range of species measured at different times of the year in their natural Mediterranean environment. As for stress detection, Penuelas et al. (1997b) showed that WI was a good indicator of water status in response to salinity. WI might be the most relevant spectroradiometrical index for decisionmaking in irrigation scheduling.

CONCLUSIONS

Physiological tools of common use in agronomical assessment of yield performance and environment adaptation of crop canopies could be of use, with some adjustments and/or calibrations, in the assessment of the performance of Irrigation Systems.

REFERENCES

- Araus, J.L., Febrero, A., Buxo, R., Camalich, M.D., Martin, D., Molina, F., Rodriguez-Ariza, M.O. and Romagosa, I. 1997. Changes in carbon isotope discrimination in grain cereals from different regions of the western Mediterranean basin during the past seven millennia. Palaeoenvironmental evidence of a differential change in aridity during the late Holocene. *Global Change Biology* 3, 107-118.
- Araus, J.L., Febrero, A., Catala, M., Molist, M., Voltas, J. and Romagosa, I. 1999. Crop water availability in early agriculture: evidence from carbon isotope discrimination of seeds from a tenth millennium BP site on the Euphrates. *Global Change Biology* 5, 201-212.

- Araus, J.L., Villegas, D., Aparicio, N., Garcia-del-Mora, IL.F., Elhani, S., Rharrabti, Y., Ferrio, J.P. and Royo, C. 2003. Environmental Factors Determining Carbon Isotope Discrimination and Yield in Durum Wheat Under Mediterranean Conditions. *Crop Science* 43, 170-180.
- Azzali, S. and Menenti, M. 1987. Irrigation water management in two Italian irrigation districts. Proc. Workshop on *Earthnet Pilot Project on Landsat-TM applications*, December 987, Frascati, Italy: 41–48.
- Baret, F. and Guyot, G. 1991. Potentials and limits of vegetation indices for LAI and APAR estimation. *Remote Sensing of Environment* 35, 161-173.
- Bellairs, S.M., Turner, N.C., Hick, P.T. and Smith, R.C.G. 1996. Plant and soil influences on estimating biomass of wheat in plant-breeding plots using field spectral radiometers. *Australian Journal of Agricultural Research* 47, 1017-1034.
- Ben Asher, J., Phene, C.J. and Kinarti, A. 1986. Canopy temperature to assess daily evapotranspiration and management of high frequency drip irrigation systems. *Agric. Water Management* 22(4): 379–390.
- Blackburn, G.A. 1998. Spectral indexes for estimating photosynthetic pigment concentrations A test using senescent tree leaves. *International Journal of Remote Sensing*, 19, 657-675.
- Bos, M.G. and Nugteren, J. 1974. *On Irrigation Efficiencies*. Publication no. 19, International Institute for Land Reclamation and Improvement (ILRI), Wageningen, the Netherlands: 138 pp.
- Bos, M.G., Murray-Rus, D.H., Merrey, D.J., Johnson, H.G. and Snellen, W.B. 1994. Methodologies for assessing performance of irrigation and drainage management, Irrigation and Drainage Systems 7: 231–261.
- Bos, M.G., Wolter, W., Drovandi, A. and Morabito, J.A. 1991. The Viejo Retamo Secondary Canal – performance evaluation case study: Mendoza, Argentina, Irrigation and Drainage Systems 5: 77–88.
- Burt, C.M., Clemmens, A.J., Strelkoff, T.S., Solomon, K.H., Bliesner, R.D., Hardy, L.A., Howell, T.A. and Eisenhauer, D.E. 1997. Irrigation performance measures: efficiency and uniformity. *Journal of Irrigation and Drainage Engineering* 123(6): 423–442.
- Carter, G.A. 1998. Reflectance wavebands and indexes for remote estimation of photosynthesis and stomatal conductance in pine canopies. *Remote Sensing of Environment* 63, 61-72.
- Chapelle, E.W., Kim, M.S. and Mc Murtery, J.E. 1992. Ratio analysis of reflectance spectra (RARS): an algorithm for the remote estimation of the concentrations of ChI a, b and carotenoids in soybean leaves. *Remote Sensing of Environment* 39, 239-247.
- Curran, P.J. 1983. Multispectral remote sensing for the estimation of green leaf area index. Philosophical *Transactions of the Royal Society of London Series A: Physical Sciences and Engineering* 309, 257-270.

- Elliott, G.A. and Regan, K.L. 1993. Use of reflectance measurements to estimate early cereal biomass production on sandplain soils. *Australian Journal of Experimental Agriculture* 33, 179-183.
- Farquhar, G.D., Ehleringer, J.R. and Hubick, K.T. 1989. Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology* 40, 503-537.
- Ferrio JP, Florit A, Vega A, Serrano L, Voltas J. (2003a). Δ13C and tree-ring width reflect different drought responses in Quercus ilex and Pinus halepensis. *Oecologia* 442, 512-518.
- Ferrio, J.P., Voltas, J. and Araus, J.L. 2003b. Use of carbon isotope composition in monitoring environmental changes. *Management of Environmental Quality* 14, 82-98.
- Field, C.B., Gamon, J.A. and Penuelas, J. 1994. Remote sensing of terrestrial photosynthesis. In *Ecophysiology of Photosynthesis*. (eds. E.D. Schulze & MM Caldwell), pp. 511-528. Springer- Verlag. Berlin Heidelberg.
- Filella, I. and Penuelas, J. 1995. The red edge position and shape as indicators of plant chlorophyll content, biomass and hydric status. *International Journal of Remote Sensing* 15, 1459-1470.
- Filella, I., Amaro, T., Araus, J.L and Penuelas, J. 1996. Relationship between photosynthetic radiation-use efficiency of barley canopies and the photochemical reflectance index (PRI) *Physiologia Plantarum* 96, 211-216.
- Gamon, J.A., Field, C.B., Goulden, M.L., Griffin, K.L., Hartley, A.E., Joel, G., Penuelas, J. and Valentini, R. 1995. Relationships between Ndvi, canopy structure, and photosynthesis in 3 californian vegetation types. *Ecological Applications* 5, 28-41.
- Gamon, J.A., Penuelas, J. and Field, C.B. 1992. A narrow-waveband spectral index that tracks diurnal changes in photosynthetic efficiency. *Remote Sensing of Environment* 41, 35-44.
- Gamon, J.A., Serrano, L. and Surfus, J. 1997. The photochemical reflectance index: an optical indicator of photosynthetic radiation-use efficiency across species, functional types and nutrient levels. *Oecologia* 112, 492-501.
- Hall, F.G., Huemmrich, K.F. and Goward, S.N. 1990. Use of narrow-band spectra to stimate the fraction of absorved photosynthetically active radiation. *Remote Sensing of Environment* 34, 273-288.
- Howell, T.A. and Eisenhauer, D.E. 1997. Irrigation performance measures: efficiency and uniformity. *J. of Irrigation and Drainage Engineering* 123(6): 423–442.
- Hubick, K.T., Gibson, A., Ehleringer, J.R., Hall, A.E. and Farquhar, G.D. 1993. Diversity in the relationship between carbon isotope discrimination and transpiration efficiency when water is limited. In *Stable isotopes and plant carbon-water relations* (eds. JR Ehleringer, AE Hall, and GD Farquhar) pp. 311-325. (Academic Press, Inc.: San Diego).
- Huete, A.R. 1988. A soil adjusted vegetation index (SAVI). *Remote Sensing of Environment* 25, 295-309.
- Jackson, R.D. 1984. Remote sensing of vegetation characteristics for farm management. *SPIE Remote Sensing* 475: 81–96.

- Jackson, R.D, Idso, S.B., Reginato, R.J. and Pinter, P.J. 1981. Canopy temperature as a crop water stress indicator. *Water Resources Research* 17: 1133–1138.
- Jackson, P., Robertson, M., Cooper, M. and Hammer, G. 1996. The role of physiological understanding in plant breeding; from a breeding perspective. *Field Crops Res.*, 49, 11-39.
- Jacquemoud, S. and Baret, F. 1990. PROSPECT: a model of leaf optical properties. *Remote Sensing of Environment* 34, 75-91.
- Jensen, M.E, 1977. Water Conservation and Irrigation Systems. *Climate Techn. Sem. Proc.*, Columbia, Missouri.
- Lambers, H., Chapin III, F.S and Pons, T.L. 1998. *Plant Physiological Ecology*. Springer-Verlag, New York.
- Levine, G. 1982. *Relative water supply: an explanatory variable for irrigation systems*, Technical Report no. 6, Cornell University, Ithaca, New York.
- Lichtenthaler, H., Gitelson, A. and Lang, M. 1996. Non-destructive determination of chlorophyll content of leaves of a green and an aurea mutant of tobacco by reflectance measurements. *Journal-of-Plant- Physiology* 148 (3-4) 483-493.
- Menenti, M., Visser, T.N.M., Morabito, J.A. and Drovandi, A. 1989. Appraisal of irrigation performance with satellite data and georeferenced information. In *Irrigation, Theory and Practice* (eds. Rydzewski, J.R. and Ward, C.F.), Proc. of the Int. Conf., Institute of Irrigation Studies, Southampton, 12–15 September 1989: 785–801. Pentech Press, London.
- Molden, D.J. 1997. Accounting for water use and productivity, SWIM Paper 1, International Water Management Institute, Colombo, Sri Lanka: 16 pp.
- Murray-Rust, D.H. and Snellen, W.B. 1993. *Irrigation system performance assessment and diagnosis*, IIMI, ILRI, IHE, International Irrigation Management Institute, Colombo, Sri Lanka: 148 pp.
- Penuelas, J. 1998. Visible and near-infrare reflectance techniques for diagnosing plant physiological status. *Trends in Plant Science* 3, 151-156.
- Penuelas, J., Filella, I. and Baret, F. 1995a. Semiempirical indices to assess carotenoids/chlorophyll a ratio from leaf spectral reflectance. *Photosynthetica* 31, 221-230.
- Penuelas, J., Filella, I, Biel, C., Serrano, L. and Save, R. 1993b. The reflectance at the 950-970 nm region as an indicator of plant water status. *International Journal of Remote Sensing* 14, 1887-1905.
- Penuelas, J., Filella, I. and Gamon, J.A. 1995b. Assessment of photosynthetic radiation-use efficiency with spectral reflectance. *New Phytologist* 131, 291-296.
- Penuelas, J., Filella, I., Lloret, P., Munoz, F. and Vilajeliu, M. 1995c. Reflectance assessment of plant mite attack on apple trees. *International Journal of Remote Sensing* 16, 2727-2733.
- Penuelas, J., Gamon, J.A., Griffinand, K.L. and Field, C.B. 1993a. Assessing type, biomass, pigment composition and photosynthetic efficiency of aquatic vegetation from spectral reflectance. *Remote Sensing of Environment* 46, 110-118.

- Penuelas, J., Isla, R., Filella, I. and Araus, J.L. 1997b. Visible and near-infrared reflectance assessment of salinity effects on barley. *Crop Science* 37, 198-202.
- Penuelas J, Pinol J, Ogaya R, Filella I. (1997a). Estimation of plant water concentration by the reflectance water index Wi (R900/R970). *International Journal of Remote Sensing*, 18, 2869-2875.
- Price, J.C. and Bausch, WC. 1995. Leaf-area index estimation from visible and near-infrared reflectance data. *Remote Sensing of Environment* 52, 55-65.
- Reynolds, M.P., Balota, M., Delgado, M.I.B., Amani, I. and Fischer, R.A. 1994. Physiological and morphological traits associated with spring wheat yield under hot, irrigated conditions. *Australian Journal of Plant Physiology* 21: 717-730.
- Richardson, A.J. and Wiegand, C.L. 1977. Distinguishing vegetation from soil background information. Photogrammetric. *Engineering and Remote Sensing* 43, 1541-1552.
- Rudorff, B.F.T. and Batista, G.T. 1990. Spectral response of wheat and its relationship to Agronomic variables in the tropical region. *Remote Sensing of Environment* 31, 53-63.
- Seckler, D., Sampath, R.K. and Raheja, S.K. 1988. An index for measuring the performance of irrigation management systems with an application. *Water Resources Bulletin* 24(4): 855–860.
- Shibayama, M., Wiegand, C.L. and Richardson, A.J. 1986. Diurnal patterns of bidirectional vegetation indices for wheat canopies. *International Journal of Remote Sensing* 7, 233-246.
- Small, L.E. and Svendsen, M. 1990. A framework for assessing irrigation performance, *Irrigation and Drainage Systems* 4: 283–312.
- Thomas, J.R. and Gausman, H.W. 1977. Leaf reflectance vs. leaf chlorophyll concentration for eight crops. *Agronomy Journal* 69, 799-802.
- Wiegand, C.L. and Richardson, A.J. 1990a. Use of spectral vegetation indices to infer leaf area, evapotranspiration and yield: I. Rationale. *Agronomy Journal* 82, 623-629.
- Wiegand, C.L., Richardson, A.J. 1990b. Use of spectral vegetation indices to infer leaf area, evapotranspiration and yield: II. Results. *Agronomy Journal* 82, 630-636.
- Wiegand, C.L., Richardson, A.J., Escobar, D.E. and Gerbermann, A.H. 1991. Vegetation indices in crop assessments. *Remote Sensing of Environment* 35, 105-119.
- Wolters, W. 1992. *Influences on the efficiency of irrigation water use*. PhD Thesis, Delft University of Technology, 150 pp.
- Young, A. and Britton, G. 1990. Carotenoids and stress. In Stress Responses in Plants: Adaptation and Acclimation Mechanisms. (eds. R.G. Alscher & J.R. Cumming) pp. 87-112. Wiley-Liss, New York.

Corresponding author: Dr. Hamid Kheyrodin, Faculty of Desert Science, Semnan University, Iran Email: <u>hkhyrodin@yahoo.com</u>