

Incredibility of Awesome Natural Antioxidants for Glamorous, Fascinating and Vivacious Demeanor

By

Bina Rani, Bhanwar Lal Jat, Murtaza Abid, Vikram S. Chauhan, M.M. Abid Ali Khan, Prem S. Bugasara, M.S. Naqvi, Ashok Kumar Kakodia, Piyush Panwar, Sukhraj Punar , and Raaz K. Maheshwari

ISSN 2319-3077 Online/Electronic

ISSN 0970-4973 Print

Journal Impact Factor: 4.275

Global Impact factor of Journal: 0.876

Scientific Journals Impact Factor: 3.285

InfoBase Impact Factor: 2.93

Index Copernicus International Value

IC Value of Journal 6.01 Poland, Europe

J. Biol. Chem. Research


Volume 32 (2) 2015 Pages No. 1062-1104

Journal of Biological and Chemical Research

An International Peer Reviewed / Refereed Journal of Life Sciences and Chemistry

**Indexed, Abstracted and Cited in various International and
National Scientific Databases**

Published by Society for Advancement of Sciences®



J. Biol. Chem. Research. Vol. 32, No. 2: 1062-1104, 2015

(An International Peer Reviewed / Refereed Journal of Life Sciences and Chemistry)

Ms 32/2/05/2015

All rights reserved

ISSN 0970-4973 (Print)**ISSN 2319-3077 (Online/Electronic)**

Prof. Bina Rani Dr. R. K. Maheshwari

[http:// www.sasjournals.com](http://www.sasjournals.com)[http:// www.jbcr.in](http://www.jbcr.in)jbiolchemres@gmail.cominfo@jbcr.in

REVIEW ARTICLE

Received: 15/10/2015

Revised: 21/11/2015

Accepted: 22/11/2015

Incredibility of Awesome Natural Antioxidants for Glamorous, Fascinating and Vivacious Demeanor

Bina Rani¹, Bhanwar Lal Jat², Murtaza Abid³, Vikram S. Chauhan⁴, M.M. Abid Ali Khan⁵, Prem S. Bugasara⁶, M.S. Naqvi⁷, Ashok Kumar Kakodia⁸, Piyush Panwar⁹, Sukhraj Punar¹⁰, and Raaz K. Maheshwari¹⁰

¹Department of Engineering Chemistry and Environmental Engineering, PCE, Jaipur, India

²Department of Botany, Dayanand College (MDSU) Ajmer, Rajasthan, India

³Department of Bio Chemistry, King George's Medical University, Lucknow, U.P., India

⁴Department of Botany, SBRM Govt P.G. College, Nagaur, Rajasthan, India

⁵Department of Botany, Shia P.G. College, Lucknow, U.P., India

⁶Department of Zoology, SBRM Govt. P.G. College, Nagaur, Rajasthan, India

⁷Department of Zoology, Shia P.G., College, Lucknow, U.P., India

⁸Department of Chemistry, SGG P.G. College, Banswara, Rajasthan, India

⁹Department of Botany and Biotechnology, University of Rajasthan, Jaipur, India

¹⁰Department of Chemistry, SBRM Govt. P.G. College, Nagaur, India

ABSTRACT

Dietary antioxidants, such as water-soluble vitamin C and phenolic compounds, as well as lipid-soluble vitamin E and carotenoids, present in vegetables contribute both to the first and second defense lines against oxidative stress. As a result, they protect cells against oxidative damage, and may therefore prevent chronic diseases, such as cancer, cardiovascular disease, and diabetes. While use of synthetic antioxidants (such as butylated hydroxytoluene and butylated hydroxyanisole) to maintain the quality of ready-to-eat food products has become commonplace, consumer concern regarding their safety has motivated the food industry to seek natural alternatives. Phenolic antioxidants can inhibit free radical formation and/or interrupt propagation of autoxidation. Fat-soluble vitamin E (α -tocopherol) and water-soluble vitamin C (L-ascorbic acid) are both effective in the appropriate matrix. Plant extracts, generally used for their flavoring characteristics, often have strong H-donating activity thus making them extremely effective antioxidants. This antioxidant activity is most often due to phenolic acids (gallic, protocatechuic, caffeic, and rosmarinic acids), phenolic diterpenes (carnosol, carnosic acid, rosmanol, and rosmadial), flavonoids (quercetin, catechin, naringenin, and kaempferol), and volatile oils (eugenol, carvacrol, thymol, and menthol).

Some plant pigments (anthocyanin and anthocyanidin) can chelate metals and donate H to oxygen radicals thus slowing oxidation via 2 mechanisms. Tea and extracts of grape seeds and skins contain catechins, epicatechins, phenolic acids, proanthocyanidins, and resveratrol, all of which contribute to their antioxidative activity. Mounting evidence shows that inflammation plays a part in many chronic diseases including heart disease, arthritis, asthma, diabetes, and even some cancers. This inflammation is enhanced by having too little ω -3 intake (such as in fish, flax, and walnuts), especially in relation to Omega-6 fatty acid intake (in oils such as soy and corn oil). In the quest to equalize the ratio of these two kinds of oils, flax seed can be a real help. The objective of this article is to provide an impending overview of clinical manifestation resulted due to overload of free radicals - and - natural antioxidants (-specially occurring in flaxseeds and soyabean), their mechanisms of action, nutritional prophylaxis and imminent therapeutic applicability. The significance of this area of research, current state of information, and possibilities of further investigation are precisely delineated.

Keywords: Immunocompetence, Phytochemicals, β -carotene, Free radicals, Oxidative stress, ROS, SOD, Environmental pollution, Flavonoids, PHGPX, SOD, H_2O_2 , Carotenoids, LDL, Flaxseed and Soyabean.


INTRODUCTION

The recent escalation in the knowledge of free radicals and ROS in biology is producing a medical revolution that promises a new age of health and disease management. It is ironic that oxygen, an element indispensable for life, under certain situations has deleterious effects on the human body. The demand for natural antioxidant is increasing due to its unquestionable advantages compared with the addition of antioxidants directly to the food. Therefore, the search for antioxidants perceived as natural, namely those that naturally occur in herbs and spices, is a field attracting great interest.




It is well known that many botanicals possess natural antioxidants with high antioxidant activity. And investigations on these were initiated based on their uses in traditional folkloric medicines. In the last decade, preventive medicine has undergone a great advance, especially in developed countries. Ingredients that make food functional are dietary fibers, vitamins, minerals, antioxidants, oligosaccharides, essential fatty acids (omega-3), lactic acid bacteria cultures, and lignins. Many of these are present in medicinal plants. Indian systems of medicine believe that complex diseases can be treated with complex combination of botanicals unlike in west, with single drugs.

Whole foods are hence used in India as functional foods rather than supplements. Some medicinal plants and dietary constituents having functional attributes are spices such as onion, garlic, mustard, red chilies, turmeric, clove, cinnamon, saffron, curry leaf, fenugreek, and ginger. Some herbs as *Bixa orellana* and vegetables like amla, wheat grass, soyabean, and *Gracinia cambogia* have antitumor effects. Research has demonstrated that nutrition plays a crucial role in the prevention of chronic diseases, as most of them can be related to diet. Functional food enters the concept of considering food not only necessary for living but also as a source of mental and physical well-being, contributing to the prevention and reduction of risk factors for several diseases or enhancing certain physiological functions.[1-4] A food can be regarded as functional if it is satisfactorily demonstrated to affect beneficially one or more target functions in the body, beyond adequate nutritional effects, in a way which is relevant to either the state of well being and health or reduction of the risk of a disease. The beneficial effects could be either maintenance or promotion of a state of well being or health and/or a reduction of risk of a pathologic process or a disease. [5-7]



Whole foods represent the simplest example of functional food. Broccoli, carrots, and tomatoes are considered functional foods because of their high contents of physiologically active components (sulforaphen, B-carotene, and lycopene, respectively). Green vegetables and spices like mustard and turmeric, used extensively in Indian cuisine, also can fall under this category.[9] “Nutraceutical” is a term coined in 1979 by Stephen De Felice. [8-10] It is defined “as a food or parts of food that provide medical or health benefits, including the prevention and treatment of disease.” Nutraceuticals may range from isolated nutrients, dietary supplements, and diets to genetically engineered “designer” food, herbal products, and processed products such as cereals, soups, and beverages. A nutraceutical is any nontoxic food extract supplement that has scientifically proven health benefits for both the treatment and prevention of disease. [11] The increasing interest in nutraceuticals reflects the fact that consumers hear about epidemiological studies indicating that a specific diet or component of the diet is associated with a lower risk for a certain disease. The major active nutraceutical ingredients in plants are flavonoids. As is typical for phenolic compounds, they can act as potent antioxidants and metal chelators. They also have long been recognized to possess anti-inflammatory, hepatoprotective, antithrombotic, antiviral, antiallergic, antimicrobial, antiviral, oestrogenic, enzyme inhibition, vascular and cytotoxic anti-tumor activity and anticarcinogenic activities. There is consistent evidence from human and animal studies that strenuous physical exercise may induce a state wherein the antioxidant defenses of several tissues are overwhelmed by excess reactive oxygen. There has recently been a remarkable increment in scientific articles dealing with oxidative stress.

Consequently, knowledge about reactive oxygen and nitrogen species metabolism; definition of markers for oxidative damage; evidence linking chronic diseases and oxidative stress; identification of flavonoids and other dietary polyphenol antioxidants present in plant foods as bioactive molecules; and data supporting the idea that health benefits associated with fruits, vegetables in the diet are probably linked to the polyphenol antioxidants they contain. In addition, more than 8,000 polyphenolic compounds have been identified in various plant species and reported to possess many useful properties. Brassica vegetables, which include different genus of cabbage, broccoli, cauliflower, Brussels sprouts, and kale, are consumed all over the world.




This paper aims to review the role of free radical-induced tissue damage and antioxidant defence mechanisms in inflammatory diseases that involve pathogenic processes similar to the periodontal diseases. There is a clearly defined and substantial role for free radicals or reactive oxygen species (ROS) in periodontitis. but little research has been performed in this area. This paper reviews the considerable data available relating ROS activity and antioxidant defence to inflammatory diseases and attempts to draw parallels with periodontitis. in an effort to stimulate more periodontal research in this important area. The recent discovery of the transcription factor nuclear factor κ B (NF- κ B) is reviewed and several potential pathways for cytokine-induced periodontal tissue damage, mediated by NF- κ B, are discussed. Emphasis is placed on cytokines that have been studied in periodontitis. principally TNF- α IL-1, IL-6. IL-8 and β -interferon. The link between cellular production of such important mediators of inflammation and the antioxidant (AO) thiols. cysteine and reduced glutathione (GSH), is discussed and it is hypothesised that NF- κ B antagonists may offer important therapeutic benefits.



Antioxidants may be present in foods as endogenous factors or may be added to preserve their lipid components from quality deterioration. Synthetic antioxidants such as butylated hydroxyanisole (BHA) (butylated hydroxytoluene (BHT propyl gallate (PG) and tert-butylhydroquinone (TBHQ) are commonly used in food formulations. However, due to safety concerns, interest in natural antioxidants has intensified. To address the demand by consumers, mixed tocopherols, herbal extracts such as those of rosemary and sage, as well as tea extracts and oils of soya and flax seeds, have been commercialized for food and nutraceutical applications. An overview of the topic is provided in this article.

Antioxidants are the nutritional equivalents and are loyal protectors and nurturers of our cells, repelling disease, and promoting good health. Antioxidants can come from healthy eating or in the form of supplements, and they include a family of naturally formed components like vitamin A, β -carotene, lycopene, vitamin E, and more. They are believed to protect cells from free radicals, harmful oxygen molecules thought to damage cells that result in cancer, atherosclerosis, Alzheimer's disease, and rheumatoid arthritis, and they may be the underlying reason why we age. In nature due to instinctive oxidation process - an apple slice turns brown; fish becomes rancid; a cut on skin is raw and inflamed. All of these result from a natural process called oxidation. [12]

It happens to all cells in nature, including the ones in your body. To help your body protect itself from the rigors of oxidation, Mother Nature provides thousands of different antioxidants in various amounts in fruits, vegetables, whole grains, nuts, and legumes. When your body needs to put up its best defense, especially true in today's environment, antioxidants are crucial to your health.



As oxygen interacts with cells of any type - an apple slice or, in your body, the cells lining your lungs or in a cut on your skin -- oxidation occurs. This produces some type of change in those cells. They may pass away, such as with rotting fruit. In the case of cut skin, dead cells are replaced in time by fresh, new cells, resulting in a healed cut. This birth and death of cells in the body goes on continuously, 24 hours a day.



It is a process that is necessary to keep the body healthy. [13-15] Oxidation is a very natural process that happens during normal cellular functions. Yet there is a downside. While the body metabolizes oxygen very efficiently, 1% or 2% of cells will get damaged in the process and turn into free radicals. "Free radicals" is a term often used to describe damaged cells that can be problematic.

They are "free" because they are missing a critical molecule, which sends them on a rampage to pair with another molecule. "These molecules will rob any molecule to quench that need. When free radicals are on the attack, they don't just kill cells to acquire their missing molecule. "If free radicals simply killed a cell, it wouldn't be so bad... the body could just regenerate another one," he says. "The problem is, free radicals often injure the cell, damaging the DNA, which creates the seed for disease." When a cell's DNA changes, the cell becomes mutated. It grows abnormally and reproduces abnormally -- and quickly. Normal cell functions produce a small percentage of free radicals, much like a car engine that emits fumes. But those free radicals are generally not a big problem. They are kept under control by antioxidants that the body produces naturally. External toxins, especially cigarette smoke and air pollution, are "free radical generators. Cigarette smoke is a huge source of free radicals. In fact, our food and water also harbor free radicals in the form of pesticides and other toxins. Drinking excessive amounts of alcohol also triggers substantial free radical production. Free radicals trigger a damaging chain reaction, and that's the crux of the problem.





Epidemiological studies and a substantial body of evidence have linked the production of free radicals with the occurrence of cardiovascular diseases, carcinogenesis, rheumatoid arthritis and degenerative processes associated with aging. Antioxidants aid in the prevention by scavenging the excess free radicals, hence preventing the formation of reactive oxygen species in the body. The use of synthetic antioxidants such as butylated hydroxytoluene, butylated hydroxyanisole, tert-butylhydroquinone and propyl gallate has been negatively perceived by consumers due to safety and health effects. Hence, there is an increasing interest in the search of natural antioxidants from plant sources. Most of the potentially harmful effects of oxygen are due to the formation and activity of a number of chemical compounds, known as ROS, which have a tendency to donate oxygen to other substances. Free radicals and antioxidants have become commonly used terms in modern discussions of disease mechanisms. Free radicals are formed naturally in the body, but their production is increased by factors such as smoking, alcohol, air pollution, infection, stress, excessive sunlight, and toxins like radiation and asbestos. A wide variety of physiological and dietary antioxidants act in concert to evade such a stress. Submaximal long-duration exercise training may augment the physiological antioxidant defenses in several tissues; however, this enhanced protection may not be sufficient to completely protect highly fit individuals from exhaustive exercise-induced oxidative stress. Regular physical activity in association with dietary habits that ensure adequate supply of a combination of appropriate antioxidants may be expected to yield desirable results.

Carotenoids get involved in an entire antioxidant network of physiological systems of the body. A number of nutrients must come together in synergy to afford effective antioxidant protection. Carotenoids are sacrificial antioxidants and help save other antioxidants by means of absorbing free radical hits. One mole of carotenoid absorb up to ~ 15 attacks, facilitating redox reaction continuing without uninterrupted mode of action. Weakness of one antioxidant moiety results collapse of entire network.

OXIDATIVE STRESS, FREE RADICALS and ANTIOXIDANTS

We are exposed to electromagnetic radiation from the environment, both natural and from man-made sources. Low-wavelength electromagnetic radiation (e. g., γ - rays) can split water in the body to generate hydroxyl radical, OH^* .

This fearsomely-reactive radical, once generated, attacks whatever it is next to. Its lifetime *in vivo* is vanishingly small because hydroxyl radical reacts at its site of formation, usually leaving behind a legacy in the form of propagating free-radical chain reactions.



Pathways of ROS generation and clearance

The body makes another oxygen radical (ie, the unpaired electron is located on oxygen), superoxide ($O_2^{\cdot -}$). Superoxide is made by adding one electron to the oxygen molecule. It is generally poorly reactive. Some superoxide is made by "accidents of chemistry", in that many molecules in the body react directly with oxygen to make superoxide. Examples include the catecholamines, tetrahydrofolates, and some constituents of mitochondrial and other electron-transport chains. Activated phagocytes (neutrophils, monocytes, macrophages, eosinophils) generate large amounts of superoxide as part of the mechanism by which foreign organisms are subsided. During chronic inflammations, this normal protective mechanism may become damaging. About 1-3% of the oxygen we breathe in is used to make superoxide. Since human beings consume a lot of oxygen. Another physiological free radical is nitric oxide (NO^*), which is made by vascular endothelium as a relaxing factor, and also by phagocytes and in the brain. Nitric oxide has many useful physiological functions, but excess NO can be toxic. Neither superoxide nor NO is highly reactive chemically, but under certain circumstances they can generate more toxic products. Free radicals are dangerous because they don't just damage one molecule. One free radical can set off a whole chain reaction. When a free radical oxidizes a fatty acid, it changes that fatty acid into a free radical, which then damages another fatty acid.

It's a very rapid chain reaction." These external attacks can overwhelm the body's natural free-radical defense system. In time, and with repeated free radical attacks that the body cannot stop, that damage can lead to a host of chronic diseases, including cancer, heart disease, Alzheimer's disease, and Parkinson's disease. Oxidative damage in skin cells is caused by cumulative sunlight. But if free radicals are in an internal organ - for example, if asbestos is in your lungs -- it stimulates free radical reactions in lung tissue. Cigarette smoke has active free radical generators. That's why stopping smoking is the biggest step anyone can take to preserving their health. In the 21st century, people need to get more antioxidants in their diet to offset all these assaults. These toxins are ubiquitous in the environment.

The oxidative burden on the body is much, much, much higher than it was 200 years ago, in dweller of a city. It's a fact of modern life, so we have to take that into consideration. Someone compensate for the effects of environmental toxins if consume multiple servings of fruits and veggies. Body simply doesn't produce enough antioxidants to do all that. What exactly do they do? Antioxidants work to stop this damaging, disease-causing chain reaction that free radicals have started. Each type of antioxidant works either to prevent the chain reaction or stop it after it's started. For example, the role of vitamin C is to stop the chain reaction before it starts. It captures the free radical and neutralizes it. Vitamin E is a chain-breaking antioxidant. Wherever it finds sit in a membrane, it breaks the chain reaction.



Flavonoids are the biggest class of antioxidants. Some 5,000 flavonoids in various foods have identified in various veggies, cereals, fruits and spices. Polyphenols are a smaller class of antioxidants, which often refer to as "phenols". We have clear science about antioxidants, that our bodies need a NADN for lack of a better term like a country needs a military system, the human body needs defense workers at all levels -- lieutenants, corporals, generals, staff sergeants - in the form of antioxidants. The body needs a mix of vitamins and minerals, such as vitamins A, C, E, and beta-carotene, to neutralize this free radical assault. We can't rely on a few blockbuster foods to do the job," says Blumberg. "You can't eat nine servings of broccoli a day and expect it to do it all. We need to eat many different foods. Each type works in different tissues of the body, in different parts of cells. Some are good at quenching some free radicals, some are better at quenching others. When you have appropriate amounts of different antioxidants, you're doing what you can to protect yourself. Multivitamins and vitamin supplements can provide the body with an antioxidant boost. Yet getting too much of some supplements, like vitamin E, can be harmful. Fruits, vegetables, whole grains, legumes, and nuts contain complex mixes of antioxidants, and therein lies the benefit of eating a variety of healthy foods. Researchers continue delving into the mysteries of fruits and vegetables, identifying the complex antioxidants they contain. Quercetin, luteolin, hesperetin, catetchin, even (-)-epigallocatechin are some of the blockbuster flavonoids in our foods.

Oxidative damage results when the critical balance between free radical generation and antioxidant defenses is unfavorable. Oxidative stress, arising as a result of an imbalance between free radical production and antioxidant defenses, is associated with damage to a wide range of molecular species including lipids, proteins, and nucleic acids. Short-term oxidative stress may occur in tissues injured by trauma, infection, heat injury, hypertoxia, toxins, and excessive exercise. These injured tissues produce increased radical generating enzymes (e.g., xanthine oxidase, lipogenase, cyclooxygenase) activation of phagocytes, release of free iron, copper ions, or a disruption of the electron transport chains of oxidative phosphorylation, producing excess ROS. The initiation, promotion, and progression of cancer, as well as the side-effects of radiation and chemotherapy, have been linked to the imbalance between ROS and the antioxidant defense system.

ROS have been implicated in the induction and complications of diabetes mellitus, age-related eye disease, and neurodegenerative diseases such as Parkinson's disease. A role of oxidative stress has been postulated in many conditions, including atherosclerosis, inflammatory condition, certain cancers, and the process of aging. Oxidative stress is now thought to make a significant contribution to all inflammatory diseases (arthritis, vasculitis, glomerulonephritis, lupus erythematosus, adult respiratory diseases syndrome), ischemic diseases (heart diseases, stroke, intestinal ischemia), hemochromatosis, AIDS (acquired immunodeficiency syndrome), emphysema, organ transplantation, gastric ulcers, hypertension and preeclampsia, neurological disorder (Alzheimer's disease, Parkinson's disease, muscular dystrophy), alcoholism, smoking-related diseases, and many others.

Heart diseases continue to be the biggest killer, responsible for about half of all the deaths. The oxidative events may affect cardiovascular diseases therefore; it has potential to provide enormous benefits to the health and lifespan. Poly unsaturated fatty acids occur as a major part of the low density lipoproteins (LDL) in blood and oxidation of these lipid components in LDL play a vital role in atherosclerosis. The 3 most important cell types in the vessel wall are endothelial cells; smooth muscle cell and macrophage can release free radical, which affect lipid peroxidation. With continued high level of oxidized lipids, blood vessel damage to the reaction process continues and can lead to generation of foam cells and plaque the symptoms of atherosclerosis. Oxidized LDL is atherogenic and is thought to be important in the formation of atherosclerosis plaques. Furthermore, oxidized LDL is cytotoxic and can directly damage endothelial cells. Antioxidants like β -carotene or vitamin E play a vital role in the prevention of various cardiovascular diseases.

Reactive oxygen and nitrogen species, such as super oxide anion, H_2O_2 , hydroxyl radical, and NO and their biological metabolites also play an important role in carcinogenesis. ROS induce DNA damage, as the reaction of free radicals with DNA includes strand break base modification and DNA protein cross-links. Numerous investigators have proposed participation of free radicals in carcinogenesis, mutation, and transformation; it is clear that their presence in biosystem could lead to mutation, transformation, and ultimately cancer. Induction of mutagenesis, the best known of the biological effect of radiation, occurs mainly through damage of DNA by the $HO\cdot$ Radical and other species are produced by the radiolysis, and also by direct radiation effect on DNA, the reaction effects on DNA. The reaction of $HO\cdot$ Radicals is mainly addition to double bond of pyrimidine bases and abstraction of H from the sugar moiety resulting in chain reaction of DNA. These effects cause cell mutagenesis and carcinogenesis lipid peroxides are also responsible for the activation of carcinogens.

Antioxidants can decrease oxidative stress induced carcinogenesis by a direct scavenging of ROS and/or by inhibiting cell proliferation secondary to the protein phosphorylation. β -carotene may be protective against cancer through its antioxidant function, because oxidative products can cause genetic damage. Thus, the photo protective properties of β -carotene may protect against UV light induced carcinogenesis.

Immunoenhancement of β -carotene may contribute to cancer protection. β -carotene may also have anticarcinogenic effect by altering the liver metabolism effects of carcinogens. Vitamin C may be helpful in preventing cancer. The possible mechanisms by which vitamin C may affect carcinogenesis include antioxidant effects, blocking of formation of nitrosamines, enhancement of the immune response, and acceleration of detoxification of liver enzymes. Vitamin E, an important antioxidant, plays a role in immunocompetence by increasing humoral antibody protection, resistance to bacterial infections, cell-mediated immunity, the T-lymphocytes tumor necrosis factor production, inhibition of mutagen formation, repair of membranes in DNA, and blocking micro cell line formation. Hence vitamin E may be useful in cancer prevention and inhibit carcinogenesis by the stimulation of the immune system.




The administration of a mixture of the above three antioxidant revealed the highest reduction in risk of developing cardiac cancer. The human body is in constant battle to keep from aging. Research suggests that free radical damage to cells leads to the pathological changes associated with aging. An increasing number of diseases or disorders, as well as aging process itself, demonstrate link either directly or indirectly to these reactive and potentially destructive molecules. The major mechanism of aging attributes to DNA or the accumulation of cellular and functional damage. Reduction of free radicals or decreasing their rate of production may delay aging. Some of the nutritional antioxidants will retard the aging process and prevent disease. Based on these studies, it appears that increased oxidative stress commonly occurs during the aging process, and antioxidant status may significantly influence the effects of oxidative damage associated with advancing age. Research suggests that free radicals have a significant influence on aging, that free radical damage can be controlled with adequate antioxidant defense, and that optimal intake of antioxidant nutrient may contribute to enhanced quality of life.

Recent research indicates that antioxidant may even positively influence life span. Proteins can be oxidatively modified in three ways: oxidative modification of specific amino acid, free radical mediated peptide cleavage, and formation of protein cross-linkage due to reaction with lipid peroxidation products. Protein containing amino acids such as methionine, cysteine, arginine, and histidine seem to be the most vulnerable to oxidation. Free radical mediated protein modification increases susceptibility to enzyme proteolysis. Oxidative damage to protein products may affect the activity of enzymes, receptors, and membrane transport. Oxidatively damaged protein products may contain very reactive groups that may contribute to damage to membrane and many cellular functions.

Peroxyl radical is usually considered to be free radical species for the oxidation of proteins. ROS can damage proteins and produce carbonyls and other amino acids modification including formation of methionine sulfoxide and protein carbonyls and other amino acids modification including formation of methionine sulfoxide and protein peroxide. Protein oxidation affects the alteration of signal transduction mechanism, enzyme activity, heat stability, and proteolysis susceptibility, which leads to aging.

Oxidative stress and oxidative modification of biomolecules are involved in a number of physiological and pathophysiological processes such as aging, atherosclerosis, inflammation and carcinogenesis, and drug toxicity. Lipid peroxidation is a free radical process involving a source of secondary free radical, which further can act as second messenger or can directly react with other biomolecule, enhancing biochemical lesions. Lipid peroxidation occurs on polysaturated fatty acid located on the cell membranes and it further proceeds with radical chain reaction.

Hydroxyl radical is thought to initiate ROS and remove hydrogen atom, thus producing lipid radical and further converted into diene conjugate. Further, by addition of oxygen it forms peroxyl radical; this highly reactive radical attacks another fatty acid forming lipid hydroperoxide and a new radical. Thus lipid peroxidation is propagated. Due to lipid peroxidation, a number of compounds are formed, for example, alkanes, malanoaldehyde, and isoprotanes. These compounds are used as markers in lipid peroxidation assay and have been verified in many diseases such as neurogenerative diseases, ischemic reperfusion injury, and diabetes. Many experiments clearly provide evidences that DNA and RNA are susceptible to oxidative damage. It has been reported that especially in aging and cancer, DNA is considered as a major target. Oxidative nucleotide as glycol, dTG, and 8-hydroxy-2-deoxyguanosine is found to be increased during oxidative damage to DNA under UV radiation or free radical damage. It has been reported that mitochondrial DNA are more susceptible to oxidative damage that have role in many diseases including cancer. It has been suggested that 8-hydroxy-2-deoxyguanosine can be used as biological marker for oxidative stress.





The term antioxidant originally was used to refer specifically to a chemical that prevented the consumption of oxygen. In the late 19th and early 20th century, extensive study was devoted to the uses of antioxidants in important industrial processes, such as the prevention of metal corrosion, the vulcanization of rubber, and the polymerization of fuels in the fouling of internal combustion engines. Early research on the role of antioxidants in biology focused on their use in preventing the oxidation of unsaturated fats, which is the cause of rancidity.

Antioxidant activity could be measured simply by placing the fat in a closed container with oxygen and measuring the rate of oxygen consumption. However, it was the identification of vitamins A, C, and E as antioxidants that revolutionized the field and led to the realization of the importance of antioxidants in the biochemistry of living organisms.

The possible mechanisms of action of antioxidants were first explored when it was recognized that a substance with antioxidative activity is likely to be one that is itself readily oxidized. Research into how vitamin E prevents the process of lipid peroxidation led to the identification of antioxidants as reducing agents that prevent oxidative reactions, often by scavenging ROS before they can damage cells. Antioxidants act as radical scavenger, hydrogen donor, electron donor, peroxide decomposer, singlet oxygen quencher, enzyme inhibitor, synergist, and metal-chelating agents. Both enzymatic and nonenzymatic antioxidants exist in the intracellular and extracellular environment to detoxify ROS.

Two principle mechanisms of action have been proposed for antioxidants. The first is a chain-breaking mechanism by which the primary antioxidant donates an electron to the free radical present in the systems. The second mechanism involves removal of ROS/reactive nitrogen species initiators (secondary antioxidants) by quenching chain-initiating catalyst. Antioxidants may exert their effect on biological systems by different mechanisms including electron donation, metal ion chelation, co-antioxidants, or by gene expression regulation.

The antioxidants acting in the defense systems act at different levels such as preventive, radical scavenging, repair and de novo, and the fourth line of defense, i.e., the adaptation.



The first line of defense is the preventive antioxidants, which suppress the formation of free radicals. Although the precise mechanism and site of radical formation *in vivo* are not well elucidated yet, the metal-induced decompositions of hydroperoxides and hydrogen peroxide must be one of the important sources. To suppress such reactions, some antioxidants reduce hydroperoxides and hydrogen peroxide beforehand to alcohols and water, respectively, without generation of free radicals and some proteins sequester metal ions. Glutathione peroxidase, glutathione-s-transferase, phospholipid hydroperoxide glutathione peroxidase (PHGPX), and peroxidase are known to decompose lipid hydroperoxides to corresponding alcohols. PHGPX is unique in that it can reduce hydroperoxides of phospholipids integrated into biomembranes. Glutathione peroxidase and catalase reduce hydrogen peroxide to water. *The second line of defense* is the antioxidants that scavenge the active radicals to suppress chain initiation and/or break the chain propagation reactions.

Various endogenous radical-scavenging antioxidants are known: some are hydrophilic and others are lipophilic. Vitamin C, uric acid, bilirubin, albumin, and thiols are hydrophilic, radical-scavenging antioxidants, while vitamin E and ubiquinol are lipophilic radical-scavenging antioxidants. Vitamin E is accepted as the most potent radical-scavenging lipophilic antioxidant. The third line of defense is the repair and *de novo* antioxidants. The proteolytic enzymes, proteinases, proteases, and peptidases, present in the cytosol and in the mitochondria of mammalian cells, recognize, degrade, and remove oxidatively modified proteins and prevent the accumulation of oxidized proteins. The DNA repair systems also play an important role in the total defense system against oxidative damage.

Various kinds of enzymes such as glycosylases and nucleases, which repair the damaged DNA, are known. There is another important function called adaptation where the signal for the production and reactions of free radicals induces formation and transport of the appropriate antioxidant to the right site.

Cells are protected against oxidative stress by an interacting network of antioxidant enzymes. Here, the superoxide released by processes such as oxidative phosphorylation is first converted to hydrogen peroxide and then further reduced to give water. This detoxification pathway is the result of multiple enzymes, with superoxide dismutases catalyzing the first step and then catalases and various peroxidases removing hydrogen peroxide. SODs are a class of closely related enzymes that catalyze the breakdown of the superoxide anion into oxygen and hydrogen peroxide. SOD enzymes are present in almost all aerobic cells and in extracellular fluids. There are three major families of superoxide dismutase, depending on the metal cofactor: Cu/Zn (which binds both copper and zinc), Fe and Mn types (which bind either Fe or Mn), and finally the Ni type which binds nickel. In higher plants, SOD isozymes have been localized in different cell compartments. Mn-SOD is present in mitochondria and peroxisomes. Fe-SOD has been found mainly in chloroplasts but has also been detected in peroxisomes, and Cu, Zn-SOD has been localized in cytosol, chloroplasts, peroxisomes, and apoplast. In humans (as in all other mammals and most chordates), 3 forms of superoxide dismutase are present. SOD1 is located in the cytoplasm, SOD2 in the mitochondria, and SOD3 is extracellular. The first is a dimer (consists of 2 units), while the others are tetramers (4 subunits). SOD1 and SOD3 contain Cu and Zn, while SOD2 has Mn in its reactive center.


Catalase is a common enzyme found in nearly all living organisms, which are exposed to oxygen, where it functions to catalyze the decomposition of hydrogen peroxide to water and oxygen. Hydrogen peroxide (H_2O_2) is a harmful by-product of many normal metabolic processes: to prevent damage, it must be quickly converted into other, less dangerous substances. To this end, catalase is frequently used by cells to rapidly catalyze the decomposition of H_2O_2 into less reactive gaseous O_2 and H_2O molecules. All known animals use catalase in every organ, with particularly high concentrations occurring in the liver. The glutathione system includes glutathione, glutathione reductase, glutathione peroxidases, and glutathione S-transferases. This system is found in animals, plants, and microorganisms. Glutathione peroxidase is an enzyme containing four selenium-cofactors that catalyze the breakdown of hydrogen peroxide and organic hydroperoxides. There are at least 4 different glutathione peroxidase isozymes in animals. Glutathione peroxidase 1 is the most abundant and is a very efficient scavenger of H_2O_2 , while glutathione peroxidase 4 is most active with lipid hydroperoxides.

The glutathione S-transferases show high activity with lipid peroxides. These enzymes are at particularly high levels in the liver and also serve in detoxification metabolism.

Ascorbic acid or "vitamin C" is a monosaccharide antioxidant found in both animals and plants. As it cannot be synthesized in humans and must be obtained from the diet, it is a vitamin. Most other animals are able to produce this compound in their bodies and do not require it in their diets. In cells, it is maintained in its reduced form by reaction with glutathione, which can be catalyzed by protein disulfide isomerase and glutaredoxins. Ascorbic acid is a reducing agent and can reduce and thereby neutralize ROS such as hydrogen peroxide. In addition to its direct antioxidant effects, ascorbic acid is also a substrate for the antioxidant enzyme ascorbate peroxidase, a function that is particularly important in stress resistance in plants. Glutathione is a cysteine-containing peptide found in most forms of aerobic life. It is not required in the diet and is instead synthesized in cells from its constituent amino acids. Glutathione has antioxidant properties since the thiol group in its cysteine moiety is a reducing agent and can be reversibly oxidized and reduced. In cells, glutathione is maintained in the reduced form by the enzyme glutathione reductase and in turn reduces other metabolites and enzyme systems as well as reacting directly with oxidants. Due to its high concentration and central role in maintaining the cell's redox state, glutathione is one of the most important cellular antioxidants. In some organisms, glutathione is replaced by other thiols, such as by mycothiol in the actinomycetes, or by trypanothione in the kinetoplastids.


Melatonin, also known chemically as N-acetyl-5-methoxytryptamine, is a naturally occurring hormone found in animals and in some other living organisms, including algae. Melatonin is a powerful antioxidant that can easily cross cell membranes and the blood-brain barrier. Unlike other antioxidants, melatonin does not undergo redox cycling, which is the ability of a molecule to undergo repeated reduction and oxidation. Melatonin, once oxidized, cannot be reduced to its former state because it forms several stable end-products upon reacting with free radicals. Therefore, it has been referred to as a terminal (or suicidal) antioxidant. Vitamin E is the collective name for a set of 8 related tocopherols and tocotrienols, which are fat-soluble vitamins with antioxidant properties. Of these, α -tocopherol has been most studied as it has the highest bioavailability, with the body preferentially absorbing and metabolizing this form. It has been claimed that the α -tocopherol form is the most important lipid-soluble antioxidant, and that it protects membranes from oxidation by reacting with lipid radicals produced in the lipid peroxidation chain reaction. This removes the free radical intermediates and prevents the propagation reaction from continuing. This reaction produces oxidized α -tocopheroxyl radicals that can be recycled back to the active reduced form through reduction by other antioxidants, such as ascorbate, retinol, or ubiquinol. Uric acid accounts for roughly half the antioxidant ability of plasma. In fact, uric acid may have substituted for ascorbate in human evolution. However, like ascorbate, uric acid can also mediate the production of active oxygen species.

If 2 free radicals meet, they can join their unpaired electrons and make a covalent bond (a shared pair of electrons). Thus superoxide and nitric oxide combine:



At physiological pH, peroxynitrite damages proteins directly, and decomposes into toxic products that include nitrogen dioxide gas (NO_2 and), hydroxyl radical, and nitronium ion (NO_2^+).

Hence at least some of the toxicity of excess nitric oxide may involve its interaction with superoxide. In addition, superoxide can react with iron and Cu- ions, eventually to make hydroxyl radical. Radical plus non-radical most molecules in the body are not radicals. Hence any reactive free radical generated is likely to react with a non-radical. When a free radical reacts with a non-radical, a free-radical chain reaction results and new radicals are formed.



Chemicals with multiple = bonds neutralizes free radicals by reacting with them.

Attack of reactive radicals on membranes or lipoproteins starts lipid peroxidation, which is particularly implicated in the development of atherosclerosis. If hydroxyl radicals are generated close to DNA, they can attack the purine and pyrimidine bases and cause mutations. For example, guanine is converted into 8-hydroxyguanine and other products. Superoxide dismutases (SOD) convert superoxide to hydrogen peroxide (H_2O_2):



FREE RADICAL GENERATION AT THE FOREFRONT

A free radical can be defined as any molecular species capable of independent existence that contains an unpaired electron in an atomic orbital. The presence of an unpaired electron results in certain common properties that are shared by most radicals. Many radicals are unstable and highly reactive.

They can either donate an electron to or accept an electron from other molecules, therefore behaving as oxidants or reductants. The most important oxygen-containing free radicals in many disease states are hydroxyl radical, superoxide anion radical, H_2O_2 oxygen singlet, hypochlorite, nitric oxide radical, and peroxynitrite radical. These are highly reactive species, capable in the nucleus, and in the membranes of cells of damaging biologically relevant molecules such as DNA, proteins, carbohydrates, and lipids. Free radicals attack important macromolecules leading to cell damage and homeostatic disruption. Targets of free radicals include all kinds of molecules in the body. Among them, lipids, nucleic acids, and proteins are the major targets. Free radicals and other ROS are derived either from normal essential metabolic processes in the human body or from external sources such as exposure to X-rays, ozone, cigarette smoking, air pollutants, and industrial chemicals. Free radical formation occurs continuously in the cells as a consequence of both enzymatic and nonenzymatic reactions. Enzymatic reactions, which serve as source of free radicals, include those involved in the respiratory chain, in phagocytosis, in prostaglandin synthesis, and in the cytochrome P-450 system. Free radicals can also be formed in nonenzymatic reactions of oxygen with organic compounds as well as those initiated by ionizing reactions.

Free radical reactions are expected to produce progressive adverse changes that accumulate with age throughout the body. Such "normal" changes with age are relatively common to all. However, superimposed on this common pattern are patterns influenced by genetics and environmental differences that modulate free radical damage.



These are manifested as diseases at certain ages determined by genetic and environmental factors. Cancer and atherosclerosis, two major causes of death, are salient “free radical” diseases. Cancer initiation and promotion is associated with chromosomal defects and oncogene activation. It is possible that endogenous free radical reactions, like those initiated by ionizing radiation, may result in tumor formation. The highly significant correlation between consumption of fats and oils and death rates from leukemia and malignant neoplasia of the breast, ovaries, and rectum among persons over 55 years may be a reflection of greater lipid peroxidation. Studies on atherosclerosis reveal the probability that the disease may be due to free radical reactions involving diet-derived lipids in the arterial wall and serum to yield peroxides and other substances. These compounds induce endothelial cell injury and produce changes in the arterial walls.


Exercise appears to increase reactive oxygen species, which can result in damage to cells. Exercise results in increased amounts of malondialdehyde in blood and pentane in breath; both serve as indirect indicators of lipid peroxidation. However, not all studies report increases; these equivocal results may be due to the large inter subject variability in response or the nonspecificity of the assays. Some studies have reported that supplementation with vitamins C and E, other antioxidants, or antioxidant mixtures can reduce symptoms or indicators of oxidative stress as a result of exercise. However, these supplements appear to have no beneficial effect on performance. Exercise training seems to reduce the oxidative stress of exercise, such that trained athletes show less evidence of lipid peroxidation for a given bout of exercise and an enhanced defense system in relation to untrained subjects. Whether the body's natural antioxidant defense system is sufficient to counteract the increase in reactive oxygen species with exercise or whether additional exogenous supplements are needed is not known, although trained athletes who received antioxidant supplements show evidence of reduced oxidative stress. Until research fully substantiates that the long-term use of antioxidants is safe and effective, the prudent recommendation for physically active individuals is to ingest a diet rich in antioxidants. An antioxidant is a molecule stable enough to donate an electron to a rampaging free radical and neutralize it, thus reducing its capacity to damage. These antioxidants delay or inhibit cellular damage mainly through their free radical scavenging property.

These low-molecular-weight antioxidants can safely interact with free radicals and terminate the chain reaction before vital molecules are damaged. Some of such antioxidants, including glutathione, ubiquinol, and uric acid, are produced during normal metabolism in the body. Other lighter antioxidants are found in the diet.

Although there are several enzymes system within the body that scavenge free radicals, the principle micronutrient (vitamins) antioxidants are vitamin E (α -tocopherol), vitamin C (ascorbic acid), and B-carotene. The body cannot manufacture these micronutrients, so they must be supplied in the diet.

ANTIOXIDANTS OF NATURAL and SYNTHETIC ORIGIN

Synthetic and natural food antioxidants are used routinely in foods and medicine especially those containing oils and fats to protect the food against oxidation. There are a number of synthetic phenolic antioxidants, butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) being prominent examples. These compounds have been widely used as antioxidants in food industry, cosmetics, and therapeutic industry.



However, some physical properties of BHT and BHA such as their high volatility and instability at elevated temperature, strict legislation on the use of synthetic food additives, carcinogenic nature of some synthetic antioxidants, and consumer preferences have shifted the attention of manufacturers from synthetic to natural antioxidants. In view of increasing risk factors of human to various deadly diseases, there has been a global trend toward the use of natural substance present in medicinal plants and dietary plants as therapeutic antioxidants. It has been reported that there is an inverse relationship between the dietary intake of antioxidant-rich food and medicinal plants and incidence of human diseases. The use of natural antioxidants in food, cosmetic, and therapeutic industry would be promising alternative for synthetic antioxidants in respect of low cost, highly compatible with dietary intake and no harmful effects inside the human body. Many antioxidant compounds, naturally occurring in plant sources have been identified as free radical or active oxygen scavengers. Attempts have been made to study the antioxidant potential of a wide variety of vegetables like potato, spinach, tomatoes, and legumes. There are several reports showing antioxidant potential of fruits. Strong antioxidants activities have been found in berries, cherries, citrus, prunes, and olives. Green and black teas have been extensively studied in the recent past for antioxidant properties since they contain up to 30% of the dry weight as phenolic compounds.

Apart from the dietary sources, Indian medicinal plants also provide antioxidants and these include (with common/ayurvedic names in brackets) *Acacia catechu* (kair), *Aegle marmelos* (Bengal quince, Bel), *Allium cepa* (Onion), *A. sativum* (Garlic, Lahasuna), *Aleo vera* (Indain aloe, Ghritkumari), *Amomum subulatum* (Greater cardamom, Bari elachi), *Andrographis paniculata* (Kiryat), *Asparagus recemosus* (Shatavari), *Azadirachta indica* (Neem, Nimba), *Bacopa monniera* (Brahmi), *Butea monosperma* (Palas, Dhak), *Camellia sinensis* (Green tea), *Cinnamomum verum* (Cinnamon), *Cinnamomum tamala* (Tejpat), *Curcma longa* (Turmeric, Haridra), *Embllica officinalis* (Inhian gooseberry, Amlaki), *Glycyrrhiza glabra* (Yashtimudhu), *Hemidesmus indicus* (Indian Sarasparilla, Anantamul), *Indigofera tinctoria*, *Mangifera indica* (Mango, Amra), *Momordica charantia* (Bitter gourd), *Murraya koenigii* (Curry leaf), *Nigella sativa* (Black cumin), *Ocimum sanctum* (Holy basil, Tusil), *Onosma echiioides* (Ratanjyot), *Picrorrhiza kurroa* (Katuka), Piper beetle, *Plumbago zeylancia* (Chitrak), *Sesamum indicum*, *Sida cordifolia*, *Spirulina fusiformis* (Alga), *Swertia decursata*, *Syzgium cumini* (Jamun), *Terminalia ariuna* (Arjun), *Terminalia bellarica* (Beheda), *Tinospora cordifolia* (Heart leaved moonseed, Guduchi), *Trigonella foenum-graecium* (Fenugreek), *Withania somifera* (Winter cherry, Ashwangandha), and *Zingiber officinalis* (Ginger).

In recent years, cereals and products by virtue of their rich nutritional composition and bioactive components which are beneficial for human health are being labeled as nutraceuticals, used in formulati8ng functional foods. One of such food is flaxseed (*Linum usitatissimum*) which comes in 2 varieties – brown and yellow or golden, with most types having similar nutritional values and equal amounts of ω -3 fatty acids. It may be tiny, but it's mighty: The flax seed carries one of the biggest nutrient payloads on the planet. And while it's not technically a grain, it has a similar vitamin and mineral profile to grains, while the amount of fiber, antioxidants, and Omega-3 fatty acids in flax leaves grains in the dust. Additionally, flax seed is very low in carbohydrates, making it ideal for people who limit their intake of starches and sugars. And its combination of healthy fat and high fiber content make it a great food for weight loss and maintenance -- many dieters have found that flax seed has been a key to keeping them feeling satisfied. Yes, flax seed is high in most of the B vitamins, Mg and Mn, but this little seed is just getting started. There are 3 additional nutrient groups which flax seed has in abundance, and each has many benefits. ω -3 fatty acids are a key force against inflammation in our bodies. Mounting evidence shows that inflammation plays a part in many chronic diseases including heart disease, arthritis, asthma, diabetes, and even some cancers. This inflammation is enhanced by having too little ω -3 intake (such as in fish, flax, and walnuts), especially in relation to Omega-6 fatty acid intake (in oils such as soy and corn oil). In the quest to equalize the ratio of these two kinds of oils, flax seed can be a real help. Most of the oil in flax seeds is alpha linolenic acid (ALA). ALA is an ω -3 that is a precursor to the fatty acids found in salmon and other fatty cold-water fish (called EPA and DHA). Because not everyone is able to easily convert ALA into EPA and (especially) DHA, it is best not to rely solely on flax for your Omega-3 intake. However, ALA also has good effects of its own, and definitely helps in the Omega 3/6 balance. One'd be hard-pressed to find a food higher in fiber -- both soluble and insolue-- than flax. This fiber is probably mainly responsible for the cholesterol-lowering effects of flax. Fiber in the diet also helps stabilize blood sugar, and, of course, promotes proper functioning of the intestines. : Flax seed is high in phytochemicals, including many antioxidants.

It is perhaps our best source of lignans, which convert in our intestines to substances that tend to balance female hormones. There is evidence that lignans may promote fertility, reduce peri-menopausal symptoms, and possibly help prevent breast cancer. In addition, lignans may help prevent Type 2 diabetes.



Flaxseed: An emerging nutraceutical possessing potent antioxidative activity

Flaxseeds (also called linseeds) are a rich source of micronutrients, dietary fiber, manganese, vitamin B1, and the essential fatty acid alpha-linolenic acid, also known as ALA or omega-3. Flaxseed is a source of healthy fat, antioxidants, and fiber; modern research has found evidence to suggest that flaxseed can also help lower the risk of diabetes cancer, and heart disease. King Charlemagne of the 8th century believed so strongly in the health benefits of flaxseed that he demanded his loyal subjects eat the seeds and passed laws to make sure of it. The Latin name for flax is *Linum usitatissimum*, which means "the most useful. This MNT Knowledge Center feature is part of a collection of articles on the health benefits of popular foods. It contains a nutritional profile on flaxseed, numerous health benefits associated with its consumption, as well as side effects. To reap the most benefits from flaxseeds, they should be bought in ground form or ground before consumption as whole flaxseeds can sometimes pass through the digestive tract undigested. There are two main types of flaxseed: golden flaxseed and brown flaxseed. Their nutritional profiles are very similar and both contain the same number of short-chain omega-3 fatty acids. The ANA American Nutrition Association highlighted the importance of this "neglected food," stating that flaxseed is not only "an excellent source of two fatty acids that are essential for human health - linoleic acid and alpha-linolenic acid," but also, "an excellent source of fiber and a good source of minerals and vitamins."¹ Flaxseed is very low in Cholesterol and Sodium.

Nutritional value of flaxseed/ 20 g (2 tbsp)

Energy - 54.7 kcal (DV = 3%)	Carbohydrates - 3.0 g (DV = 1%)
Sugars - 0.2 g	Dietary fiber - 2.8 g (DV = 11%)
Fat - 4.3 g (DV = 7%)	Saturated fat - 0.4 g (DV = 2%)
Monounsaturated fat - 0.8 g	Polyunsaturated fat - 2.9 g
Protein - 1.9 g (DV = 4%)	Thiamine (vit B1) - 0.2mg (DV = 11%)
Riboflavin 0.0mg	Niacin (vit. B3) - 0.3mg (DV = 2%)
Pantothenic acid (B5) - 0.1mg (DV = 1%)	Vitamin B6 - 0.0 mg
Folate - 8.9 mcg (DV = 2%)	Vitamin C - 0.1 mg (DV = 0%)
Calcium - 26.1mg (DV = 3%)	Iron - 0.6mg (DV = 3%)
Magnesium - 40.2 mg (DV = 10%)	Phosphorus - 65.8mg (DV = 7%)
Potassium - 83.3 mg (DV = 2%)	Zinc - 0.4mg (DV = 3%)

Source: USDA Nutrient Database

Lignans are one of the major classes of phytoestrogens. They are estrogen-like chemical compounds with antioxidant qualities, able to scavenge free radicals in the body. Flaxseed is considered to be one of the best sources of lignans (0.3 g/ 100g). A study, published in the journal *Critical Reviews in Clinical Laboratory Sciences*, concluded that "lignan-rich diets may be beneficial, particularly if consumed for life." *Flaxseeds (also called linseeds) are a rich source of micronutrients, dietary fiber, manganese, vitamin B1, and the essential fatty acid alpha-linolenic acid, also known as ALA or omega-3.* Flaxseed is rich in both soluble fiber (dissolves in water) and insoluble fiber (does not dissolve in water). According to Mayo Clinic³, "soluble fiber dissolves with water and creates a gel-like substance that helps to lower blood cholesterol and glucose levels." Whereas insoluble fiber "absorbs water, which adds bulk to your digestive tract and helps to move things through quickly." These are considered to be "good fats" that are beneficial for the heart. These essential acids are only obtainable by eating the right foods; the human body is not able to produce them. The therapeutic and beneficial properties of consuming flaxseed are not yet completely understood, and many claims still lack "high-quality" studies to back them up. However, emerging research suggests that flaxseed might indeed be the wonder food many people claim it to be.

Consuming flaxseed may help protect against prostate, colon, and breast cancers. Flaxseed is thought to prevent the growth of cancerous cells because its omega-3 fatty acids disrupt malignant cells from clinging onto other body cells. In addition, the lignans in flaxseed have antiangiogenic properties - they stop tumors from forming new blood vessels. One US study presented at the 43rd annual meeting of the American Society of Clinical Oncology (ASCO) revealed that consuming flaxseed can stop prostate cancer tumors from growing. Dr Wendy Demark-Wahnefried, lead investigator of the study said that the team was "excited that this study showed that flaxseed is safe and associated with a protective effect on prostate cancer." Researchers at the ISU's NWRC found that cholesterol levels lowered among men who included flaxseed in their diet. People who can't take something like Lipitor, this could at least give you some of that cholesterol-lowering benefit." A study published in the Journal of the Society for Integrative Oncology suggests that a dietary intake of flaxseed can decrease the risk of hot flashes among postmenopausal women. Not only does flaxseed seem to alleviate hot flashes, but it appears to have overall health and psychological benefits as well. There is strong evidence to suggest that consuming flaxseed every day improves glycemic control in obese men and women with pre-diabetes. . A diet of flaxseed may protect skin tissue from being damaged by radiation, revealed researchers at the PSMPU Perelman School of Medicine at the University of Pennsylvania. The investigators concluded that their "study demonstrates that dietary flaxseed, already known for its strong antioxidant and anti-inflammatory properties, works as both a mitigator and protector against radiation pneumonopathy. [115-125]


Soybean and Healthcare

Soybean (*Glycine max* L), assumes a predominant position in solving the problem of food shortage the world over. A native of China, soybean has been cultivated for food well over 13,000 years. The Chinese name for soybean means 'greater bean'. Like other beans, soybeans grow in pods, containing edible seeds. While we most often think of them as being green, the seeds can also be yellow, brown or black. Today soybeans are grown all over the world.

This plant was introduced in most countries as a source of oil food and protein for livestock but now it is commercially grown for many food and industrial purposes. About 70% of the total production goes for oil extraction and rest for seed purposes (10%) and direct food uses (20%). The oil so obtained is refined and used for culinary purposes. It is also used as an important ingredient for industrial products such as paints, plastics, lubricants and bio-fuels. The main by-product of the oil industry, namely lecithin (phospholipid) finds commercial application as a nutritional supplement and emulsifier. Other by-product includes hulls, which are used in animal feeds and as a source of fiber. The meal primarily used as a source of protein for poultry, piggery, livestock, aquaculture, etc. Soy meal has more than 50% edible grade protein, which can also be diverted for food uses. However the meal from the solvent extraction plants must be made edible grade and devoid of the residual solvents, which may cause various physiological disorders in humans.

The International Standard Organisation (ISO) recommends 50ppm of residual hexane while Bureau of Indian Standards (BIS) allows 170ppm of such residual solvents. In an innovative process, developed by the International Soybean Center (INTSOY) at the University of Illinois, the soya bean oil is extracted primarily through extrusion. The meal is devoid of the solvent and also contains low profile of fat. It may be used for direct food uses through either supplementation or fortification with traditional foods.

Soy foods contain all nine essential amino acids. The protein digestibility corrected amino acid score (PDCAAS) is 1.0, which is equivalent to animal protein. Soybeans have a number of nutritional advantages over other food legumes. Soybean derives about 35 to 38% of its calories from protein compared to approximately 20 to 30% in other legumes. It contains on an average 40% protein that is much higher when compared to other legumes.



The values for other legumes are: chickpea, 4.9-29.6%; peas, 21.2-32.9%; cowpea, 20.9-38.5; pigeon pea, 18.8-28.5%; green gram, 20.8-33.1%; lentil, 20.4-30.5% and lathyrus, 22.7-29.6%. The quality of soya proteins is almost at par with egg or milk proteins, which are ideal with reference to their essential amino acid make-up. Like other pulses, soybeans are deficient in sulphur-containing amino acids such as cystine and methionine. Since it has more lysine than cereals, its blending with them makes the product well balanced. It is therefore suggested to blend soybean with cereals, millets and other pulses at different proportions as per our body requirement. Approximately 40% of the calories in soybeans are derived from fats. Soybean has exceptionally large quantities of fat. It has on an average 20% oil. The oil is hypocholesterolemic.

The oil content is much higher than other pulses such as black gram, 1.64%; pigeon pea, 2.19%; cowpea, 2.05%; chick pea, 4.99%; lentil, 1.17%; lathyrus, 1.0% and green gram, 2.14%. The quality of oil is normally judged by its fatty acid composition. More the unsaturated fatty acids better the quality of oil. It contains about 78% of unsaturated fatty acids. Out of them linoleic and linolenic constitute 58%.

They are called polyunsaturated fatty acids (PUFA). The iodine value is in the range of 125-135. The oil remains liquid over a relatively wide range. The oil can be hydrogenated selectively for blending with semi solid or liquid oils. Naturally occurring antioxidants/tocopherols are present and are not completely removed during processing. However, soya oil has certain disadvantage like high phosphate content (2%), which must be removed by processing. The oil also contains 7-8% linolenic acid, which is responsible for flavour and odor reversion. It contains about 20-30% carbohydrates. Mostly they are galacto oligosaccharides such as raffinose, stachyose and verbascose. It has little starch. The carbohydrate content is much lower than the other legumes where the major portion is starch such as black gram, 56.5-63.7%; chickpea, 60.1-61.7; green gram, 53.3-61.2; pigeon pea, 57.3-58.7% and lentil, 59.7-61.0%. It also has considerable amounts of calcium (226mg); phosphorus (546mg); iron (8.5mg); iron (8.5mg); magnesium (236mg); copper (2.4mg); and sodium (27.9mg) per 100g of beans. All whole, unprocessed plant foods contain dietary fiber. One serving of soybeans provides approximately 8 grams of dietary fiber. However, many soya foods are processed in ways that decrease their fiber content significantly. Tofu and soya milk, two of the more popular soya foods, contain very little fiber. Soya foods that utilize the whole bean such as tempeh, soya flour and textured soya protein, are high in fiber. About 30% of the fiber in soya foods is soluble fiber. It is a rich source of vitamin A (426mg); thiamine (73mg); riboflavin (39mg) and niacin (3.2mg)/100g beans. Soya foods are the richest source of isoflavones. These are phyto-serms (selective estrogen receptor modulators). They have some estrogen-like qualities and have non-hormonal properties as well. The two primary isoflavones in soybeans are genistein and diadzein and their glycosides. They contribute to many protective effects. Consumption, of soya protein provides health benefits that may help prevent or treat certain chronic diseases. Soy foods and other soy based dairy analogs can serve as a balanced and remedial substitute of dairy milk for lactose intolerant persons. This condition arises mainly due to lack of beta galactosidase, the enzyme responsible for the hydrolysis of lactose in the intestine. The lactose is in turn degraded by the colonic bacteria into acid and carbon dioxide causing gastric discomfort such as flatulence, bloating, belching and diarrhoea. Since soybean has no lactose in it, the products prepared from soybean, namely, soy paneer and other soymilk analogues can serve as an ideal substitute of regular milk. There is some evidence that soy foods may help with sugar control in diabetics. Soy may also help lower risk of some of the complications of diabetes, such as kidney disease. Soybeans have a very low glycemic index and are valuable in a diabetic diet. Blood sugar control may also be improved by choosing carbohydrates that are high in soluble fiber. It helps in the slow absorption of the sugars. In kidney disease, a soy-based diet may be preferable to the traditional low protein diet from decreasing the renal damage. Soy provides high quality protein without stimulating hyperfiltration and proteinuria. Soy protein lowers the total and LDL cholesterol levels. Soy foods are excellent choice for a heart-healthy diet. Soy oil provides the plant-derived omega-3 fatty acid, ALA, while fish oil contains the marine-derived omega-3 fatty acids, EPA and DHA.

These omega-3 fatty acids improve heart function by providing greater variability between beats, therefore reducing the risk of arrhythmia and/ or sudden death. Soy in the diets will have significant reduction in both diastolic and systolic blood pressure. Not only the total blood cholesterol is significantly lowered, the level of HDL (good) cholesterol are also significantly increased. Soy protein can reduce high blood cholesterol levels by 10 to 15% - enough to cut the chances of a heart attack by upto 30%. Soy protein inhibits cholesterol oxidation. Oxidised cholesterol is cholesterol that has undergone structural changes because of exposure to oxygen. Only oxidised cholesterol can damage arteries. The hormonal changes that occur during menopause can cause a variety of symptoms and increase risk for heart disease and osteoporosis. During perimenopause women experience fluctuations in estrogen levels. This can cause hot flashes, night sweats, insomnia, vaginal dryness or headaches. Hormonal Replacement Therapy (HRT) is commonly prescribed to help prevent the negative health effects of menopause. However, many women do not want to take HRT because of the possible increased risk for breast cancer. Soy foods, which contain isoflavones, may decrease the health risk associated with menopause. They also lower the rates of osteoporosis and heart disease leading to longer expectancy.: Soy foods contain anticarcinogens that may prove protective. They lower the incidences of breast, prostate and colon cancers. Soy foods fit in the formulation of a health promoting diet. The fiber in soybeans also provides preventive therapy for several other conditions. high-fiber soybeans may be able to help reduce the risk of colon cancer. As a matter of fact, in areas of the world where soybeans are eaten regularly, rates of colon cancer, as well as some other cancers, including breast cancer, tend to be low. Soybean fiber may also be able to reduce the symptoms of diarrhoea or constipation in suffers o irritable bowel syndrome. Today's soy formulas are equivalent to cows' milk formulas in digestibility, nutrition profile and acceptability. In healthy infants, soy formulas promote normal growth, nutritional status and bone mineralization. Substitution of soy protein for a protein for a protein source of animal origin can result in reduction of calorie. Soy protein concentrate and soy protein isolate are reported to have 330 cal/100g. The soy-based diet thus lowers the incidences of obesity. Active isoflavone compounds found in soy, specifically, genistein help us stay lean by producing fewer and smaller fat cells. **Oxidative** damage is often the result of high levels of unhealthy free radicals in the body. Free radicals are unstable or highly reactive moieties that cause unhealthy effects to our body's metabolism. Getting enough antioxidants in our diet to promote healthier tissues is essential to reduce the unwanted effects of these free radicals.

RESEARCH TRIUMPH

An extensive review of the literature on the role of oxidative stress in influencing assisted reproduction and its outcome is described by Agarwal et al. (2006). Free radicals or reactive oxygen species mediate their action through many of the proinflammatory cytokines and this mechanism has been proposed as a common underlying factor for endometriosis, ovarian cancer, polycystic ovary disease, and various other pathologies affecting the female reproductive process, as highlighted in this review. Oxidative stress, sperm DNA damage, and apoptosis have been implicated in male infertility. Elevated reactive oxygen species levels correlate with the poor fertility outcomes seen in the assisted reproductive technology setting.

Oxidative stress has been implicated in male and female infertility, including fetal dysmorphogenesis, abortions, and intrauterine growth restriction. Accurate evaluation of seminal oxidative stress by standardized assays may help in the diagnosis and management of male infertility. There is evidence in the literature on the beneficial effects of oral antioxidant supplementation in male infertility. Current ongoing trials will provide answers on the safety and effectiveness of antioxidants in improving maternal and fetal outcomes. Further studies need to be conducted to determine if antioxidant supplementation will prevent fetal developmental defects in high-risk pregnancy with diabetes.


Asthma is a chronic disease characterized by inflammation of the airways (Buse and Lemanske2001), a complex disorder characterized by variable and recurring symptoms, airflow obstruction, bronchial hyperresponsiveness, and an underlying inflammation. The interaction of these features determines the clinical manifestations and severity of asthma, and it has been reported as a disease of increasing prevalence. The pathogenesis of asthma is unknown but imbalances between oxidants and antioxidants are believed to play a fundamental role. One key component of the oxidant-antioxidant hypothesis centers on the huge burden of oxidants derived from inflammatory cell infiltration into the lung. The eosinophil, in particular, is implicated as a major source of oxidative injury, including protein nitration (MacPherson et al. 2011).

Rank	Fruits and Veggies	Serving Size	Antioxidant Capacity/ Serving
1	Small red beans, dried	1/2 cup	13727
2	Wild blueberries	1 cup	13427
3	Red kidney beans, dried	1/2 cup	13259
4	Pinto beans	1/2 cup	11864
5	Blueberries, cultivated	1 cup	9019
6	Cranberries	1 cup	8983
7	Artichoke hearts, cooked	1 cup	7904
8	Blackberries	1 cup	7701
9	Dried prunes	1/2 cup	7291
10	Raspberries	1 cup	6058
11	Strawberries	1 cup	5938
12	Red delicious apple	One	5900
13	Granny Smith apple	One	5381
14	Pecans	1 ounce	5095
15	Sweet cherries	1 cup	4873
16	Black plum	One	4844
17	Russet potato, cooked	One	4649
18	Black beans	1/2 cup	4181
19	Plum	1	4118
20	Gala apple	1	3903

Dysfunctional mitochondria in lung cells are another potential source of oxidants. Mitochondrial injury to airway epithelium occurs in murine models of allergic asthma (Aguilera-Aguirre et al. 2009; Mabalirajan et al. 2008).

THE MOST COMPREHENSIVE STUDY OF THE ANTIOXIDANT CONTENT OF COMMON FOODS

There is evidence to support its role in human asthma as well, including increased oxidative injury to mitochondrial epithelial cell superoxide dismutase (SOD) (Comhair et al. 2005), enhanced mitochondrial proliferation in bronchial smooth muscle (Trian et al. 2007), and mutations in mitochondrial DNA (Reddy 2011). Overall, this oxidative burden, generated by both inflammatory and lung cells, can overwhelm antioxidant defense to cause oxidant stress during asthma.



This stress can alter or inactivate the function of essential proteins, lipids and nucleic acids culminating in severe cell injury, dysfunction and death. Among many unknown and complicated mechanisms, involvement of airways inflammation with an oxidant/antioxidant imbalance such as reactive oxygen species (ROS) can lead to lung injury as a result of direct oxidative damage to epithelial cells and cells shedding.

As inflammation is often associated with an increased generation of reactive oxygen species (ROS), it is rational to surmise that an oxidant stress could be mechanistically important in asthma. ROS have been shown to be associated with the pathogenesis of asthma by inducing bronchial hyper reactivity as well as directly stimulating histamine release from mast cells and mucus secretion from airway epithelial cells (Ryszard 2000).

The great external surface area (1–2) m² of the human airway epithelium plus its direct contact with the environment, makes the respiratory tract a major target for oxidative injury from inhaled oxidants such as cigarette smoke, ozone, hyperoxia, nitrogen and sulphur oxides and other airborne pollutants. It has been well recognized that biological systems are capable of forming highly reactive moieties, both free radicals and non-radicals named reactive oxygen species (ROS) and reactive nitrogen species (RNS). Free radicals can especially be generated in a wide variety of chemical and biological systems, including the formation of plastics, the ageing of paints, the combustion of fuels and in the human body. In living organisms, the levels of free radicals and other 'reactive species' are controlled by a complex web of antioxidant defences, which minimize (but do not completely prevent) oxidative damage to biomolecules (Roberfroid and Calderon 1995; Gaston et al. 1994; Halliwell (2005).

These biologically active species serve in cell signaling as messenger molecules of the autocrine or paracrine system (Saran and Bors 1989; Suzuki et al. 1997) and also in host defense, (biocidal effects against microbial and tumor cells) (Babior 1978) but their excessive production may result in tissue injury and inflammation (Halliwell et al. 1992; Gutteridge and Halliwell 1994). Reportedly, any excessive production of oxidants is kept to a minimum by a well coordinated and efficient endogenous antioxidant defense mechanism. It has been proposed that a deficit in the precise balance between exposure to oxidants and endogenous antioxidants results in oxidative stress which appears to be involved in the pathogenesis of a growing number of diseases, including lung pathologies such as respiratory distress syndrome, asthma, idiopathic and iatrogenic pulmonary fibrosis, cystic fibrosis, HIV-associated lung disease, lung cancer and other pulmonary diseases and conditions (Clement and Housset 1996; Barnes 1995) . As excessive ROS levels damage lipids, proteins and nucleic acids through oxidation and thus are associated with various diseases, such as atherosclerosis, arthritis, neurodegenerative disorders, and cancers , a regular supplement of antioxidants can assist the endogenous defense systems to counterbalance the harmful effects of excessive ROS (Balsano and Alisi 2009; Kaur and Geetha 2006).

Another source of natural antioxidants is naturally occurring from traditional Chinese medicines sources, have been identified as free radical or active oxygen scavengers (Duh 1998; Pan et al.2007). Natural antioxidants can protect the human body from free radicals and retard the progress of many chronic diseases as well as retard lipid oxidative rancidity in foods or medicinal materials (Kang et al. 2008). Among the many mentioned sources of naturally occurring antioxidant, Seahorse (*Hippocampus kuda* Bleeker) has been well known for its special medicinal composition. According to Zhong-Ji Qian et al., the methanol extracts of seahorse contained high amount of phenolic compounds and those extracts exhibited good antioxidant activity by effectively scavenging various free radicals such as DPPH radicals, hydroxyl radicals, superoxide anion radicals, alkyl radicals, and reducing the ferric to ferrous ion in different antioxidant systems.

In a search for treatment that might stop the recurrent attacks of breathlessness and wheezing to make it more susceptible to at least, providing relief for asthmatic patients, and if possible to treat the asthmatic disease, a method has emerged that seems to be extremely useful for application of natural sources of antioxidants for treatment of asthma. The method was using finely extracted powders from the seahorse (*Hippocampus kuda*) and *Rhizoma Homalomenae* (with a ratio of 1: 1 w/w) in honey to form into pill of 500mg. All the hand-rolled pills were dried in an oven at 55°C until the moisture content of the pill was consistent.

In this paper, successful application of extracted powders from the seahorse (*Hippocampus kuda*) and *Rhizoma Homalomenae* together with honey in the form of medical pill for treatment of asthma is reported.

Phenolic antioxidants such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT) and monotertiary butylhydroquinone (TBHQ) are hindered phenols with antimicrobial activity. The antimicrobial activity of phenolic antioxidants appears to depend on the presence of a hydroxyl group on the molecule, the lipid solubility of the compound and the degree of steric hindrance.



The use of phenolic antioxidants in food products is regulated by federal agencies. In general, these compounds are permitted in concentrations up to 200 ppm, based on the fat or oil content of the food product. Certain food products have special regulations. The antimicrobial activity of phenolic antioxidants has been studied in meat and its products, poultry and its products, milk and its products, seafood, rice, applesauce and food ingredients.

The antimicrobial activity of phenolic antioxidants is modified by at least 10 factors such as microbial species/strain, stressed microorganisms, type and concentration of phenolic antioxidants, concentration of microbial challenge, combination of phenolic antioxidants, combination of phenolic antioxidants with other antimicrobials, combination of phenolic antioxidants with temperature and food additives, food components, carriers of phenolic antioxidants and the mode of addition of phenolic antioxidants. The antimicrobial activity of phenolic antioxidants in foods has been examined against growth and by-products of bacteria (gram positive and negative, spore and nonspore formers, spoilage and pathogenic), molds and yeasts. The concentration of phenolic antioxidants that had antimicrobial activity in food products was in the range of 30–10,000 ppm. The mechanism of inhibition by phenolic antioxidants has been found to affect the function and composition of the cellular membrane, the synthesis of DNA, RNA, protein and lipid, and the function of the mitochondrion.

Radical scavenging capacity and antioxidant activity of an E vitamer fraction from rice bran were investigated. The E vitamer fraction was prepared by a liquid-liquid extraction method. The free radical scavenging capacity of the E vitamer fraction was measured by the 1,1-diphenyl-2-picrylhydrazyl radical (DPPH) method at the concentration range of 2.5 to 640 ppm, whereas the antioxidant activities were measured by both the reducing power and ferric thiocyanate (FTC) methods at different concentrations (0, 2.5, 10, 40, and 160 ppm). Radical scavenging capacity of the fraction was effective up to 160 ppm, then remained stable. Also, at a concentration of 160 ppm, it was highly effective in inhibiting linoleic acid peroxidation.



Similarly, superoxide scavenging and antioxidant activities of the E vitamer fraction were compared with those of the synthetic phenol compounds. Results showed that with a concentration of the E vitamer fraction at 160 ppm, the antioxidant activity was comparable to both butylated hydroxytoluene (BHT, 160 ppm) and butylated hydroxyanisole (BHA, 160 ppm). No significant differences ($P > 0.05$) were found among them.

The E vitamer fraction may be a good substitute for the synthetic phenol antioxidants currently used in industry because the former is more natural and comparatively effective in its radical scavenging capacity and antioxidant activity.

During 16 h heating at 180 °C, the oxidative stability (OS) of virgin olive oil (VOO) as affected by the same concentrations (200 ppm) of tertiary-butylhydroquinone (TBHQ) and unsaponifiable matters of bene kernel (UKO) and hull (UHO) oils in terms of the inhibitory effect on the formation of conjugated diene hydroperoxides (OS_{CDV}) and off-flavor carbonyl compounds (OS_{CV}) was investigated. TBHQ was not able to considerably increase the OS_{CDV} (7.51) of the VOO (7.2) and showed no synergistic effect with indigenous antioxidative compounds of the VOO (IOV) in this respect. However, it could significantly improve the OS_{CV} (from 2.49 to 4.52), which was mainly due to its synergism with the IOV. The UKO increased considerably the OS_{CDV} (to 11.8), and its OS_{CV} (4.22) was nearly the same as that of TBHQ. The IOV still had marked contributions to the prevention of VOO oxidation but the majority of stabilizing effect was related to the UKO and its synergism with the IOV. The OS_{CDV} in presence of the UHO was less than that of the VOO (5.96), although it significantly increased the OS_{CV} (to 5.2), mainly due to the stabilizing effect of UHO and its synergism with the IOV.

In vitro reducing power tests of tea polyphenols TBHQ, and monoglyceride citrate (MGC) revealed that, at certain concentrations, Tp + TBHQ displayed a strong reductive capacity. Moreover, DPPH radical-scavenging activity tests showed that Tp + TBHQ was effective ($p < 0.05$) in quenching DPPH radicals. The effectiveness of Tp + TBHQ towards the radical-scavenging activity was also found to be increased when their concentrations were increased.

The oxidation stability of FO was investigated as a function of time at high temperature, in the absence and presence of antioxidants; the PV and *p*-anisidine value of FO were determined as indicators of oxidation at 60°C during 20 days. The results revealed that $Tp_{200} + TBHQ_{100}$ (subscripts refer to amounts in mg/kg) had noticeable reduction effects on the oxidation (PV and *p*-anisidine) of FO compared with $TBHQ_{200}$ regarding the PV, and when compared with the mixture of $TBHQ_{100} + MGC_{100}$ regarding the *p*-anisidine value. However, according to Rancimat tests, the ternary mixture ($Tp_{200} + TBHQ_{100} + MGC_{100}$) was found to be more effective in controlling FO oxidation. The induction periods for FO containing $Tp_{200} + TBHQ_{100} + MGC_{100}$, $TBHQ_{100} + MGC_{100}$, and $Tp_{200} + TBHQ_{100}$ were 8.305, 6.905, and 6.8020 h, respectively, whereby no significant differences were observed between $TBHQ_{100} + MGC_{100}$ and $Tp_{200} + TBHQ_{100}$ ($p < 0.025$).

Practical applications: Flaxseed oil is an unstable vegetable oil due to its high content of PUFA, which are mainly composed of ALA. One effective way to ensure a high quality of lipids and lipid-containing products, and to prolong their storage time, is to stabilize them by addition of suitable antioxidants. The stabilization of FO will help to increase both its shelf life and to improve its use in different applications. According to recent studies, FO contains many components that are beneficial for human health, especially ALA which can contribute to increasing the DHA and EPA content in the body. FO is believed to provide benefits in the prevention and treatment of heart diseases, arthritis, inflammatory, and autoimmune diseases.


The vitamin E family consists of eight isomers known as alpha-, beta-, gamma-, and delta-tocopherols and alpha-, beta-, gamma-, and delta-tocotrienols. Numerous studies focused on the health benefits of these isomers have been performed since the discovery of vitamin E in 1922. Recent discoveries on the potential therapeutic applications of tocotrienols have revolutionized vitamin E research. Nevertheless, despite the abundance of literature, only 1% of vitamin E research has been conducted on tocotrienols. Many new advances suggest that the use of tocotrienols for health improvement or therapeutic purposes is promising. Although the mechanisms of action of tocotrienols in certain disease conditions have been explored, more detailed investigations into the fundamentals of the health-promoting effects of these molecules must be elucidated before they can be recommended for health improvement or for the treatment or prevention of disease. Furthermore, many of the studies on the effects of tocotrienols have been carried out using cell lines and animal models. The effects in humans must be well established before tocotrienols are used as therapeutic agents in various disease conditions, hence the need for more evidence-based human clinical trials.

The antifungal effect of BHA, BHT, TBHQ and PC alone or in combination on three toxigenic strains of aspergilli (NRRL 2999, NRRL 4123, NRRL 5835) and three nontoxigenic strains of aspergilli (NRRL 5521, NRRL 5917, NRRL 5918) was examined in a solid medium and in salami. BHT and PG (0.001, 0.005, 0.01, 0.02g per plate) did not inhibit growth, sporulation, and toxigenesis of all six cultures. Aflatoxin production by toxigenic aspergilli (B_1 , B_2 , G_1 and G_2) in the presence of BHA, TBHQ, and a combination of BHA and TBHQ was reduced significantly ($P < 0.05$). In salami BHA, TBHQ alone or in combination at 100 ppm significantly ($P < 0.05$) decreased the aflatoxin production by aspergilli when compared to control samples. A combination of BHA and TBHQ showed synergistic inhibition in both studies (solid medium and salami studies).




Effects of synthetic phenolic antioxidants (BHA, BHT, and TBHQ) on the MB sensitized photooxidation of linoleic acid as compared with that of α -tocopherol have been studied. Their antioxidative mechanism was studied by both ESR spectroscopy in a TMPD- MB system and spectroscopic analysis of rubrene oxidation induced by a chemical source of singlet oxygen.

Total singlet oxygen quenching rate constants ($k_{ox-Q}+k_q$) were determined using a steady state kinetic equation. TBHQ showed the strongest protective activity against the MB sensitized photooxidation of linoleic acid, followed by BHA and BHT.



TBHQ (1×10^{-3} M) exhibited 86.5% and 71.4% inhibition of peroxide and conjugated diene formations, respectively, in linoleic acid photooxidation after 60-min light illumination. The protective activity of TBHQ against the photosensitized oxidation of linoleic acid was almost comparable to that of α -tocopherol. The data obtained from ESR and rubrene oxidation studies clearly showed the strong singlet oxygen quenching ability of TBHQ.

The $k_{ox-Q}+k_q$ of BHA, BHT, and TBHQ were determined to be 3.37×10^7 , 4.26×10^6 , and $1.67 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$, respectively. The $k_{ox-Q}+k_q$ of TBHQ was within the same order of magnitude of that of α -tocopherol, a known efficient singlet oxygen quencher. There was a high negative correlation ($r^2=0.991$) between $\log (= k_{ox-Q}+k_q)$ and reported oxidation potentials for the synthetic antioxidants, indicating their charge-transfer mechanism for singlet oxygen quenching. This is the 1st report on the kinetic study on $k_{ox-Q}+k_q$ of TBHQ in methanol as compared with other commonly used commercial synthetic antioxidants and α -tocopherol.



Current research suggests that wine contains substances that may reduce the mortality rate from coronary diseases. The oxidation of LDL is thought to be a key step in the development of atherosclerosis. Phenolic fractions of a Petite Syrah wine were evaluated for their antioxidant activity in inhibiting LDL oxidation *in vitro*. The more active fractions contained components of the catechin family. The catechin oligomers and the procyanidin dimers (B_2 , B_3 , B_4 , B_6 , B_8) and trimers (C_1 , C_2) were extracted, isolated and purified from grapes seeds. These compounds were tested for their inhibition of LDL oxidation, along with other monomeric wine phenolics. The procyanidin dimers B_2 and B_8 , and trimer C_1 , and the monomers catechin, epicatechin and myricetin had the highest antioxidant activity. The procyanidin dimers B_3 , B_4 and C_2 and the monomers gallic acid, quercetin, caffeic acid, and rutin, and a group of compounds that included the dimer B_6 , ellagic acid, sinapic acid, cyanidin had lower antioxidant activity and α -tocopherol had the least activity. Thus, the numerous phenolic compounds found in wine are potent antioxidants in inhibiting LDL oxidation *in vitro*.

In recent years, the concept of cancer chemoprevention has matured greatly. Significant reversal or suppression of premalignancy in several sites by chemopreventive agents appears achievable. This article summarizes experimental data on chemopreventive effects of tea polyphenols in different tumor bioassay systems. Tea (*Camellia sinensis*) is cultivated in about 30 countries, and is the most widely consumed beverage in the world. Three main commercial tea varieties—green, black, and oolong—are usually consumed, but most experimental studies demonstrating the antimutagenic and anticarcinogenic effects of tea have been conducted with water extract of green tea, or GTP (polyphenolic fraction isolated from green tea). The majority of these studies have been conducted in a mouse skin tumor model system where tea is fed either as water extract through drinking water, or as purified GTP. GTP has been shown to exhibit antimutagenic activity *in vitro*, and inhibit carcinogen—as well as UV-induced skin carcinogenesis *in vivo*. Tea consumption has also been shown to afford protection against chemical carcinogen-induced stomach, lung, esophagus, duodenum, pancreas, liver, breast, and colon carcinogenesis in specific bioassay models. Several epicatechin derivatives (polyphenols) present in green tea have been shown to possess anticarcinogenic activity; the most active is (–)-epigallocatechin-3-gallate, which is also the major constituent of GTP.

The mechanisms of tea's broad cancer chemopreventive effects are not completely understood. Several theories have been put forward, including inhibition of UV- and tumor promoter-induced ornithine decarboxylase, cyclo-oxygenase, and lipoxygenase activities, antioxidant and free radical scavenging activity; enhancement of antioxidant (glutathione peroxidase, catalase, and quinone reductase) and phase II (glutathione-S-transferase) enzyme activities; inhibition of lipid peroxidation, and anti-inflammatory activity. These properties of tea polyphenols make them effective chemopreventive agents against the initiation, promotion, and progression stages of multistage carcinogenesis.

To test the feasibility of dry milling oats (*Avena sativa* L.) to concentrate antioxidant activity and phenolic antioxidants, groats were pearled for 5 to 180 s. These treatments removed <1 to 15% of the weight. The material obtained from short pearling times was mostly bran. Longer pearling times increased the amount of starchy endosperm in the pearlings. Antioxidant activity of 80% ethanol extracts, measured by β -carotene bleaching and by reduction of the free radical, 2,2-diphenyl-1-picrylhydrazyl, was highest in the short-pearling-time fractions and decreased as more endosperm tissue was included. Likewise, there was a decreasing concentration of total phenolics, determined colorimetrically, and of several simple phenolic acids, determined by high performance liquid chromatography, as more material was pearled from the groats. In contrast, concentrations of avenanthramides were not correlated with pearling time, indicating that they were more uniformly distributed in the groats.

Natural antioxidants have recently gained increased interest because of the belief that natural food ingredients are better and safer than synthetic ones. The review presents the results on stabilisation of the main edible oils with different types of natural antioxidants. Sources of natural antioxidants are spices, herbs, teas, oils, seeds, cereals, cocoa shell, grains, fruits, vegetables, enzymes, proteins. Researchers concentrate on ascorbic acid, tocopherols and carotenoids as well as on plant extracts containing various individual antioxidants such as flavonoids (quercetin, kaemferol, myricetin), catechins or phenols (carnosol, rosmanol, rosmaridiphenol) and phenolic acids (carnosic acid, rosmarinic acid). Ascorbyl palmitate is regarded as a 'natural' antioxidant because it is hydrolysed in the body to ascorbic and palmitic acids. Among the herbs of the *Lamiaceae* family, rosemary has been more extensively studied and its extracts are the first marketed natural antioxidants. Oregano, which belongs to the same family, has gained the interest of many research groups as a potent antioxidant in lipid systems. The review concerns the following main topics: stabilisation of oil with individual natural antioxidants, interaction of antioxidants with synergists, stabilisation of oil with extracts or dry materials from different plant sources (e.g. herbs and spices), stabilisation at frying temperatures and in emulsions.

Six phenolic antioxidative compounds [5-caffeoylquinic acid (chlorogenic acid), 3,5-dicaffeoylquinic acid, quercetin 3-galactoside, quercetin 3-glucoside, quercetin 3-(6-malonylglucoside), and quercetin 3-(6-malonylgalactoside) (tentative)] were identified from the leaves of *Corchorus olitorius* L. (moroheiya) by NMR and FAB-MS.

The contents of these phenolic compounds, ascorbic acid, and α -tocopherol in *Corchorus olitorius* leaves were determined, and their antioxidative activities were measured using the radical generator-initiated peroxidation of linoleic acid. The results obtained showed that 5-caffeoylquinic acid was a predominant phenolic antioxidant in *Corchorus olitorius* leaves.

ROS, associated with depletion of antioxidant defense. In the present work we evaluated the level of lipid peroxidation and antioxidant components in blood of sportsmen under resting conditions and compared the data obtained with those in age- and sex-matched sedentary controls. A significant increase was noted in the levels of TBARS and conjugated dienes while a decrease was observed in ascorbic acid and glutathione levels in sportsmen. α -Tocopherol was unaltered in plasma of sportsmen as compared to controls. The activity of superoxide dismutase was increased (52 %) and glutathione peroxidase was decreased (43 %) in the erythrocytes of sportsmen compared to controls. Basal glutathione levels were negatively correlated with conjugated dienes and maximal oxygen uptake (VO_{2max}) of the subjects. Dietary supplementation with antioxidant vitamins has been shown to be beneficial in combating oxidative stress without enhancing performance while exogenous glutathione was found to influence the endurance capacity of athletes. Such studies demonstrate the critical role played by glutathione and suggest that intervention trials should include a mixture of antioxidants rather than a single antioxidant.

CONCLUSION

Free radicals damage contributes to the etiology of many chronic health problems such as cardiovascular and inflammatory disease, cataract, and cancer. Antioxidants prevent free radical induced tissue damage by preventing the formation of radicals, scavenging them, or by promoting their decomposition. Synthetic antioxidants are recently reported to be dangerous to human health.



Thus the search for effective, nontoxic natural compounds with antioxidative activity has been intensified in recent years. In addition to endogenous antioxidant defense systems, consumption of dietary and plant-derived antioxidants appears to be a suitable alternative.

Dietary and other components of plants form a major source of antioxidants. The traditional Indian diet, spices, and medicinal plants are rich sources of natural antioxidants; higher intake of foods with functional attributes including high level of antioxidants in antioxidants in functional foods is one strategy that is gaining importance.

Newer approaches utilizing collaborative research and modern technology in combination with established traditional health principles will yield dividends in near future in improving health, especially among people who do not have access to the use of costlier western systems of medicine. One of the most important steps you can take to prevent heart disease, cancer, Alzheimer's, and Parkinson's disease is to eat plenty of antioxidant-rich foods. Antioxidants are bounteous in plant foods, particularly those that have bright colours. Because there are many different types of antioxidants that can protect your tissues from different types of damage, it is best to eat a wide range of antioxidant-rich foods. How much you benefit from the antioxidants found in the foods you eat depends on how well you breakdown and absorb these foods. One of the best ways of making sure that you are getting plenty of antioxidants in your diet is to strive to eat lots of fresh vegetables. If you just don't have the time to eat a large green salad every day, consider buying a good juicer and drinking a fresh vegetable juice on a daily basis. Another more convenient option is to use a high quality super green food product. It is best to limit the amount of sweet fruits that you eat according to your dental health and blood sugar and insulin levels. When washing and preparing vegetables and fruits, be sure to wash non-organic varieties with extra care to help remove pesticide residues. This is especially important for vegetables and fruits that are known to be heavily contaminated with pesticides. The highest ranked foods in four major categories are as follows - Fruits: blueberries, cranberries, and blackberries; Vegetables: beans, and surprisingly, russet potatoes; nuts: pecans, walnuts, and hazelnutsp; Spices: cinnamon, oregano, and ground cloves.

ABBREVIATIONS

TBARS (thiobarbituric acid reactive substances); ROS (reactive oxygen species); GTP (green tea polyphenols); NF- κ B (transcription factor nuclear factor κ B); LDL (low-density lipoprotein); ROS (reactive oxygen species); AO (antioxidant); cysteine and reduced glutathione (GSH (glutathione); BHS (butylated hydroxyanisole); BHA (butylated hydroxytoluene); PG (propyl gallate); TBHQ (tert-butylhydroquinone); ISU's NWRC (Iowa State University's Nutrition and Wellness Research Center); PSMPU (Perelman School of Medicine, University of Pennsylvania); ANA (American Nutrition Association); ISO (The International Standard Organisation); BIS (Bureau of India Standards) INTSOY, UI (International Soybean Center, University of Illinois); PDCAAQS (protein digestibility corrected amino acid score); SOD (Superoxide dismutases); ASCO (American Society of Clinical Oncology); HRT (Hormonal Replacement Therapy); VOO (virgin olive oil); tert-butylhydroquinone (TBHQ), and MGC (monoglyceride citrate); DPPH (1,1-diphenyl-2-picrylhydrazyl); FO (flaxseed oil); PV (peroxide value); ALA (α -linolenic acid); BHA (butylated hydroxyanisole); BHT (butylated hydroxytoluene); TBHQ (tertiary butylhydroquinone); PC (propyl gallate); TMPD (2,2,6,6-tetramethylpiperidone); MB (methylene blue); LDL (low-density lipoprotein); ROS (reactive oxygen species); RNS (reactive nitrogen species); NSDN (Natural Antioxidant Defense Network).

REFERENCES

1. Duh, P.D. 1998. Antioxidant activity of burdock (*Arctium lappa* Linné): Its scavenging effect on free-radical and active oxygen. *J Am Oil Chem Soc*, 5:455-461.
2. Ghasemzadeh, A. and Ghasemzadeh, N. 2011. Flavonoids and phenolic acids: Role and biochemical activity in plants and human. *J Med Plants Res* 5(31): 6697-6703.
3. Gutteridge, J.M.C. and Halliwell, B. 1994. Antioxidants in nutrition, health and disease. Oxford: Oxford University Press.
4. Halliwell, B., Gutteridge, J.M.C. and Cross, C.E. 1992. Free radicals, antioxidants and human disease: where are we now ? *J Lab Clin Med*, 119: 598-620.
5. Halliwell, B. 2005. Free radicals and other reactive species in disease. In: eLS. Chichester: John Wiley.
6. Kaur, I.P. and Geetha, T. 2006. Screening methods for antioxidants-a review. *Mini Rev Med Chem*, 6(3):305-312.
7. Aruoma, O.I. 1994. Nutrition and health aspects of free radicals and antioxidants. *Food Chem Toxicol.*, 32:671-83.
8. Freeman, B.A., and Crapo, J.D. 1982. Biology of disease: Free radicals and tissue injury. *Lab Invest*, 47:412-26.
9. Halliwell, B. 1995. How to characterize an antioxidant- An update. *Biochem Soc Symp.* 61:73-101.
10. Maheshwari, R.K., Garg, A. and Singh, P. 2012. Red wine and its salutary inherent succession. *Bulletin of Environment, Pharmacology and Life Sciences*, 1 (7)01 - 03.
11. Maheshwari, R.K. and Karuna Mittal 2014. Soy based diet: a mesmerizing triumph for wellness. *Pharma Tutor Magazine* 2 (2): 117-123.
12. Maheshwari, R.K., Mohan, L. and Upadhyay, B. 2014. Garcinia cambogia: an awesome superfruit for healthcare. *Journal of Pharmaceutical and Biomedical Analysis Letters* 2 (1): 80-84.
13. Singh, P., Dhawan, N.G. and Maheshwari, R.K. 2014. Herbs and spices: to embellish therapeutically for healthcare. *Pharma Tutor* 2 (1): 98-107. 231.
14. German, J. 1999. Food processing and lipid oxidation. *Adv Exp Med Biol.* 459:23-50.
15. Knight, J. 1998. Free radicals: Their history and current status in aging and disease. *Ann Clin Lab Sci*; 28:331-46.
16. Sen, C.K. 1995. Oxidants and antioxidants in exercise. *Journal of Applied Physiology*, 79: 675-686.
17. Toan, N.V. and Hanh, T.T. 2013. Improved treatment of Asthma by using natural sources of antioxidants Springer Plus; 2:278 doi:10.1186/2193-1801-2-278.
18. Rebecca, S.Y. Wong, R.S.Y. and Radhakrishnan, A.K. 2012. Inhibition of *In Vitro* Human LDL Oxidation by Phenolic Antioxidants from Grapes and Wines. *Nutrition Reviews*, 70 (9):483-490. doi:10.1111/j.1753-4887.2012.00512.x
19. Teissedre, P.L., Frankel, E.N., Waterhouse, A.L.O. and Peleg H., Germa 1996. *Journal of the Science of Food and Agriculture* 70 (1): 55-61. doi: 10.1002/(SICI)1097-0010(199601)70:1<55::AID-JSFA471>3.0.CO;2-X.
20. Chapple, I.L.C. 2005. Reactive oxygen species and antioxidants in inflammatory diseases. *Journal of Clinical Periodontology* 24 (5): 287-296. doi: 10.1111/j.1600-051X.1997.tb00760.x

21. Maheshwari, R.K., Yadav, R.K., Malhotra, J. Dhawan, N.G., Mohan, L., Rajnee, Jat, B.L., Upadhayay, B. and Rani, B. 2014. Fascinating nutritional, prophylactic, therapeutic and socio-economic reconcile attributable to drum stick tree (*Moringa oleifera* Lam.). *Global Journal of Medical Research* XIV (B-I): 11-21.
22. Yanishlieva, N.V. and Marinova, E.M. 2010. Stabilization of edible oils with natural antioxidants. *Pharmacognosy Review* 4(8):118-126. doi: 10.1002/1438-9312(200111)103:11<752::AID-EJLT752>3.0.CO;2-0.
23. Halliwell, B. 1994. Free radicals, antioxidants, and human disease: curiosity, cause, or consequence ? *The Lancet* 344 (8924): 721–724.
24. Brewer, M.S. 2011. Natural antioxidants: sources, compounds, mechanisms of action, and potential applications. *Comprehensive Reviews in Food Science and Food Safety* 10 (4): 221-247. DOI: 10.1111/j.1541-4337.2011.00156.x
25. Shahidi, F. 2000. Antioxidants in food and food antioxidants LWT Food Science and Technology 4 (3): 158-163. LWT DOI: 10.1002/1521-3803 (20000501)44:3<158::AID-FOOD158>3.0.CO;2-L
26. Agarwal, A., Gupta, S. and Sikka, S. 2006. The role of free radicals and antioxidants in reproduction. *Current Opinion in Obstetrics and Gynecology* 18 (3) 325-332. Doi:10.1097/01.geo.0000193003.58158.4e
27. Santosh, K., Katiyar, S.K. and Mukhtar, H. 1997. *Journal of Cellular Biochemistry* 67 (27 Suppl.):59-67.doi: 10.1002/(SICI)1097-4644(1997)27+<59::AID-JCB11>3.0.CO;2-G
28. Priscilla, M. Clarkson, M.P. and Thompson, H.S. 2000. Antioxidants: what role do they play in physical activity and health ? *American Journal of Clinical Nutrition* 72 (2): 637s-646s
29. Rani, B., Jat, B.L., Pandey, M.K., Malhotra, J., Mohan, L., Bugasara, P.S. and Maheshwari, R.K. 2014. Dietetic corollary of *Momordica charantia* for vivaciousness and sanguinity. *International Journal of Pharmaceutical Research and Bioscience* 3 (2):141-154.
30. Maheshwari, R.K., Rani, B., Rajnee, Parihar S. and Sharma, M. 2013. Bitter gourd with astounding therapeutic worth. *International Journal of Medicine and Pharmaceutical Research* 1 (1): 135-138.
31. Maheshwari, R.K., Chauhan, A.K., Gupta, A. and Sharma, S. 2013. Cinnamon: an imperative spice for human comfort. *International Journal of Pharmaceutical Research and Bio-science* 2 (50): 131-145.
32. Adlercreutz, H. 2007. "Lignans and human health." *Critical Reviews in Clinical Laboratory Sciences*. 44 (5-6):483-525.
33. Andrea, M. Hutchins, Blakely D. Brown, Stephen C. Cunnane, Stephanie G. Domitrovich, Earle R. Adams and Courtney E. Bobowiec. 2013. "Daily flaxseed consumption improves glycemic control in obese men and women with pre-diabetes: a randomized study". *Nutrition Research*. 3 (5):367-375.
34. Rani, B., Singh, U., Sharma, R., Gupta, A., Dhawan, N.G. Sharma, A.K., Sharma, S., and Maheshwari, R. K. 2013. *Prosopis Cineraria* (L) Druce: A desert tree to brace livelihood in Rajasthan. *Asian Journal of Pharmaceutical Research and Health Care* 5 (2): 58-64.
35. Maheshwari, R.K., Parmar, V. and Joseph, L. 2013. Latent Therapeutic gains of beet root. *World Journal of Pharmaceutical Res.*, 2 (4): 804-820.
36. Rani, B., Maheshwari, R.K., Sharma, S., Parihar, S. and Singh, U. 2013. Meraculous therapeutic potential of Indian Gooseberries. *International Journal of Medicine and Pharmaceutical Research*, 1 (2): 218-225.

37. Maheshwari, R.K., Rani, B. and Parihar, S. 2013. Versatile virtues of ginger. *International Journal of Medicine and Pharmaceutical Research*, 1 (1): 135-143.
38. Maheshwari, R.K. and Rani, B. 2013. Intrinsic benefits and therapeutic applications of rose water. *Universal Journal of Pharmacy*, 2 (2): 42-46.
39. Maheshwari, R.K., Rani, B. and Parihar, S. 2013. Vigour and vitality from *Camellia sinensis* plant. . *Universal Journal of Pharmacy* 2 (2); 4-7.
40. Rani, B., Maheshwari, R.K. and Rajnee. 2013. Healthy course to livelihood with citrus fruits. *Universal Journal of Pharmacy* 2 (3): 17-24.
41. Maheshwari, R.K., Rani, B. and Parihar, S. 2013. An overview on nutritional consequence of fruits. *Universal Journal of Pharmacy* 2 (3): 52-56.
42. Rani, B., Bhati, I., Dhawan, N.G., Rajnee, Sharma S., Tyagi, S.N. and Maheshwari, R.K. 2013. Dietary antioxidants to prop up vigour and vitality: an overview. *Journal of Drug Discovery and Therapeutics* 1 (7): 106-122.
43. Tyagi, S.N., Patel, C., Dhruv, M., Ishita, M., Gupta, A.K., Usman, M.R.M., Nimbiwal, B., and Maheshwari, R.K. 2013. Benefits of *Apium graveolens* (Celery herb). *Journal of Drug Discovery and Therapeutics* 1 (5) 36-38.
44. Maheshwari, R.K., Rani, B., Dhawan, N.G. and Singh, U. 2013. Phytochemical potentiality of broccoli, tomato and garlic. *International Journal of Current Trends in Pharmaceutical Research* 1 (2): 81-87.
45. Maheshwari, R.K., Singh, U., Dhawan, N.G., Bhati, I. and Rani, B. 2013. Therapeutic pragmatism of grape fruit. *International Journal of Chemistry and Pharmaceutical Sciences* 1 (3): 187-192.
46. Maheshwari, R.K. and Kumar, M. 2013. Hazardous impacts of electromagnetic radiations on human health. *Bulletin of Environment, Pharmacology and Life Sciences* 2 (3): 66-67.
47. Asha, K.K., Suseela, M. and Lakshmaman, P.T. 2012. Flavonoids and phenolic compounds in two mangrove species and their antioxidant property. *Indian J Geo Mar Sci* 41(3):259-264.
48. Ghasemzadeh, A. and Ghasemzadeh, N. 2011. Flavonoids and phenolic acids: Role and biochemical activity in plants and human. *J Med Plants Res* 5 (31):6697-6703.
49. Gutteridge, J.M.C. and Halliwell, B. 1994. Antioxidants in nutrition, health and disease. Oxford: Oxford University Press.
50. Halliwell, B., Gutteridge, J.M.C. and Cross, C.E. 1992. Free radicals, antioxidants and human disease: where are we now ? *J Lab Clin Med* 119:598-620.
51. Kaur, I.P. and Geetha, T. 2006. Screening methods for antioxidants-a review. *Mini Rev Med Chem* 6(3):305-312.
52. Suzuki, Y.J., Forman, H.J. and Sevanian, A. 1997. Oxidants as stimulators of signal transduction. *Free Rad Biol Med* 22:269-85.
53. Urguiaga, I. and Leighton, F. 2000. Plant polyphenol antioxidants and oxidative stress. *Biol Res* 33:55-64.
54. Yen, G.C. and Chen, H.Y. 1995. Antioxidant Activity of Various Tea Extracts in Relation to Their Antimutagenicity *J. Agric. Food Chem* 43:27-32.
55. Zheng, W. and Wang, S.Y. 2001. Antioxidant activity and phenolic compounds in selected herbs. *J Agric Food Chem* 49:5165-5170.

56. Aruoma, O.I. 2003. Methodological consideration for characterization for potential antioxidant actions of bioactive components in plants foods. *Mutat Res.*; 532:9–20.
57. Maheshwari, R.K., Malhotra, J., Mohan, L. and Upadhyay, B. 2014. Exotic pepino: a shrub for prophylactic consequence and nutritional regime. *International Journal of Pharma Research and Health Sciences* 2 (1): 42-48.
58. Rani, B., Singh, U. and Maheshwari, R.K., 2013. Role of green chemistry for human comfort and economic intensification. *Unique Journal of Chemistry*, 1 (1): 1-15.
59. Maheshwari, R.K., Yadav, R.K., Malhotra, J., Dhawan, N.G., Mohan, L., Rajnee, Jat, B.L., Upadhyay, B., and Rani, B. 2014. Fascinating nutritional, prophylactic, therapeutic and socio-economic reconcile attributable to drum stick tree (*Moringa oleifera* Lam.). *Global Journal of Medical Research* XIV (B-I): 11-21.
60. Rani, B., Jat, B.L., Pandey, M.K., Malhotra, J., Mohan, L., Bugasara, P.S. and Maheshwari, R.K. (2014). Dietetic corollary of *Momordica charantia* for vivaciousness and sanguinity. *International Journal of Pharmaceutical Research and Bioscience* 3 (2):141-154.
61. Maheshwari, R.K., Rani, B., Rajnee, Parihar S. and Sharma, M. 2013. Bitter gourd with astounding therapeutic worth. *International Journal of Medicine and Pharmaceutical Research* 1 (1): 135-138.
62. Maheshwari, R.K. and Rani, B. 2014. Efficacy of ultrasound technology in food industry. *Food Marketing and Technology* 5 (4): 38-39.
63. Maheshwari, R.K., Chauhan, A.K., Mohan, L. and Maheshwari, M. 2014. Spice up for scrumptious tang, cologne and well being. *Journal of global Biosciences* 3 (1): 304-313.
64. Maheshwari, R.K. 2014. Are pesticides ostentatious on fruits and veggies. *Biocomplex* 2 (1): 72-74.
65. Maheshwari, R.K. and Kumar, M. 2013. Nutritional importance of kakadu plum. *The Science Observer* 1 (2): 11-12.
66. Maheshwari, R.K. and Kumar, M. 2013. The unique persona of Spirulina. *The Science Observer* 1 (3): 6-8.
67. Rani, B., Singh, U. and Maheshwari, R.K. 2013. Natural antioxidants and their intrinsic worth for wellbeing. *Global Journal of Medical Research* XIII (B-VII):54-69.
68. Rani, B., Jat, B.L., Pandey, M.K., Malhotra, J., Mohan, L., Bugasara, P.S. and Maheshwari, R.K. 2014. Dietetic corollary of *Momordica charantia* for vivaciousness and sanguinity. *International Journal of Pharmaceutical Research and Bioscience* 3 (2):141-154.
69. Maheshwari, R.K., Rani, B., Rajnee, Parihar, S. and Sharma, M. 2013. Bitter gourd with astounding therapeutic worth. *International Journal of Medicine and Pharmaceutical Research* 1 (1): 135-138.
70. Maheshwari, R.K. and Rani, B. 2014. Efficacy of ultrasound technology in food industry. *Food Marketing and Technology* 5 (4): 38-39.
71. Maheshwari, R.K. and Joshi, V. 2013. Intensifying apprehension for food protection. *The Science Observer*, 1 (10): 08-09.
72. Sharma, S. and Maheshwari, R.K. 2013. Pesticide in our food. *Agrobios Newsletter*, XI (08):86.
73. Sharma, S., Maheshwari, R.K. and Chaudhari, V. 2013. Dos and don'ts for wellbeing. *Agrobios Newsletter*, XI (08): 94.
74. Chaudhary, V., Yadav, R.K. and Maheshwari, R.K. 2013. Quit smoking to survive longer. *Agrobios Newsletter*, XI (10): 91.

75. Joshi, V., Dhawan, N.G. and Maheshwari, R.K. 2013. GMOs: not a solution to world hunger. *Agrobios Newslette*, XII (4); 10-11.
76. Mohan, L. and Maheshwari, R.K. 2013. Sustainability of organic farming for future perspectives. *Agrobios Newsletter*, XII (6): 41-42.
77. Maheshwari, R.K. 2013. Bio bin composting. *Agrobios Newsletter*, XII (6): 42.
78. Maheshwari, R.K., Chauhan, A.K., Gupta, A. and Sharma, S. 2013. Cinnamon: an imperative spice for human comfort. *International Journal of Pharmaceutical Research and Bio-science* 2 (50): 131-145.
79. Adlercreutz, H. 2007. "Lignans and human health." *Critical Reviews in Clinical Laboratory Sciences*. 44 (5-6):483-525.
80. Andrea, M. Hutchins, Blakely D. Brown, Stephen C. Cunnane, Stephanie G. Domitrovich, Earle R. Adams and Courtney E. Bobowiec. 2013. "Daily flaxseed consumption improves glycemic control in obese men and women with pre-diabetes: a randomized study". *Nutrition Research*. 3 (5): 367-375.
81. Rani, B., Singh, U., Sharma, R., Gupta, A., Dhawan, N.G. Sharma, A.K., Sharma, S., and Maheshwari, R. K. 2013. *Prosopis Cineraria* (L) Druce: A desert tree to brace livelihood in Rajasthan. *Asian Journal of Pharmaceutical Research and Health Care* 5 (2): 58-64.
82. Maheshwari, R.K., Yadav, R.K., Malhotra, J., Dhawan, N.G., Mohan, L., Rajnee, Jat, B.L., Upadhayay, B. and Rani B. (2014). Fascinating nutritional, prophylactic, therapeutic and socio-economic reconcile attributable to drum stick tree (*Moringa oleifera* Lam.). *Global Journal of Medical Research* XIV (B-I): 11-21.
83. Maheshwari, R.K., Parmar, V. and Joseph, L. 2013. Latent Therapeutic gains of beet root. *World Journal of Pharmaceutical Research*, 2 (4): 804-820.
84. Rani, B., Maheshwari, R.K., Sharma, S., Parihar, S., Singh, U. 2013. Meraculous therapeutic potential of Indian Gooseberries. *International Journal of Medicine and Pharmaceutical Research*, 1 (2): 218-225.
85. Maheshwari, R.K., Rani, B. and Parihar, S. 2013. Versatile virtues of ginger. *International Journal of Medicine and Pharmaceutical Research*, 1 (1): 135-143.
86. Maheshwari, R.K. and Rani, B. 2013. Intrinsic benefits and therapeutic applications of rose water. *Universal Journal of Pharmacy* 2 (2): 42-46.
87. Maheshwari, R.K., Rani, B. and Parihar, S. 2013. Vigour and vitality from *Camellia sinensis* plant. *Universal Journal of Pharmacy* 2 (2); 4-7.
88. Rani, B., Maheshwari, R.K. and Rajnee 2013. Healthy course to livelihood with citrus fruits. *Universal Journal of Pharmacy* 2 (3): 17-24.
89. Maheshwari, R.K., Rani, B. and Parihar, S. 2013. An overview on nutritional consequence of fruits. *Universal Journal of Pharmacy* 2 (3): 52-56.
90. Maheshwari, R.K., Mohan, L., Rani, B., Tyagi, S.N., Chauhan, K., Bhati, I., Rajnee and Sharma, S. 2013. Detrimental impacts of toxic ingredients found in colas. *Journal of Drug Discovery and Therapeutics* 1 (7): 34-41.
91. Rani, B., Bhati, I., Dhawan, N.G., Rajnee, Sharma, S., Tyagi, S.N. and Maheshwari, R.K. 2013. Dietary antioxidants to prop up vigour and vitality: an overview. *Journal of Drug Discovery and Therapeutics* 1 (7): 106-122.
92. Tyagi, S.N., Patel, C., Dhruv, M., Ishita, M., Gupta, A.K., Usman, M.R.M., Nimbiwal, B., and Maheshwari, R.K. 2013. Benefits of *Apium graveolens* (Celery herb). *Journal of Drug Discovery and Therapeutics* 1 (5) 36-38.

93. Maheshwari, R.K., Rani, B., Dhawan, N.G. and Singh, U. 2013. Phytochemical potentiality of broccoli, tomato and garlic. *International Journal of Current Trends in Pharmaceutical Research* 1 (2): 81-87.
94. Maheshwari, R.K., Singh, U., Dhawan, N.G., Bhati, I. and Rani, B. 2013. Therapeutic pragmatism of grape fruit. *International Journal of Chemistry and Pharmaceutical Sciences* 1 (3): 187-192.
95. Maheshwari, R.K., Rani, B., Verma, D.M. and Maheshwari, R.K. 2013. Indigenous probiotics and health benefits. *Bulletin of Environment, Pharmacology and Life Sciences* 2 (1): 3-87.
96. Maheshwari, R.K. and Kumar, M. 2013. Hazardous impacts of electromagnetic radiations on human health. *Bulletin of Environment, Pharmacology and Life Sciences* 2 (3): 66-67.
97. Maheshwari, R.K. and Rani, B. 2013. Augmentation of enzymatic antioxidants *in vitro*. *Bulletin of Environment, Pharmacology and Life Sciences* 2 (5): 101-102.
98. Maheshwari, R.K., Kumar, M. and Maheshwari, M. 2013. Sangria: a drink for fitness and cheerfulness. *International Archive of Applied Sciences and Technology* 4 (3):76-36.
99. Balakrishnan, S.D. and Anuradha, C.V. 1998. Exercise, depletion of antioxidants and antioxidant manipulation. *Cell Biochemistry and Function* 16(4):269–275. doi: 10.1002/(SICI)1099-0844 (1998120)16:4<269::AID-CBF797>3.0.CO;2-B.
100. Beckman, K.B. and Ames, B.N. 1998. The free radical theory of ageing matures. *Physiology Review* 78 (2):547-581.
101. Eberlein-Konig, B., Placzek, M. and Przybilla, B. 1998. Protective effect against sunburn of combined systematic ascorbic acid (vitamin C) and d- α -tocopherol. *Journal of American Academy of Dermatology* 38 (1):45-48.
102. Asha, K.K., Suseela, M. and Lakshmanan, P.T. 2012. Flavonoids and phenolic compounds in two mangrove species and their antioxidant property. *Indian J Geo Mar Sci* 41(3):259-264.
103. Babior, B.M. 1978. Oxygen dependent microbial killing by phagocytes. *N Engl J Med* 298:658-9.
104. Balsano, C. and Alisi, A. 2009. Antioxidant effects of natural bioactive compounds. *Curr Pharm Des* 15:3063-3073.
105. Barnes, P.J. 1995. Nitric oxide and airway disease. *Ann Med* 27:389-93.
106. Buse, W.W. and Lemanske, R.F. Jr 2001. *Asthma*. *N Engl J Med* 344:350-362.
107. Molyneux, P. 2004. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating an antioxidant activity. *Songklanakarin J Sci Technol* 26(2):211-219.
108. Natella, F., Nardini, M., Felice, M.D. and Scaccini, C. 1999. Benzoic and cinnamic acid derivatives as antioxidants: structure activity relation. *J Agric Food Chem* 47:1453-1459.
109. Pandey, K.B. and Rizvi, S.I. 2009. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev* 2(5):270-278.
110. Saran, M. and Bors, M. 1989. Oxygen radicals acting as chemical messenger: a hypothesis. *Free Rad Res Commun* 7(3–6):213-20.
111. Suzuki, Y.J., Forman, H.J. and Sevanian, A. 1997. Oxidants as stimulators of signal transduction. *Free Rad Biol Med* 22:269-85.
112. Urguiga, I. and Leighton, F. 2000. Plant polyphenol antioxidants and oxidative stress. *Biol Res* 33:55-64.

113. Yen, G.C. and Chen, H.Y. 1995. Antioxidant Activity of Various Tea Extracts in Relation to Their Antimutagenicity *J. Agric. Food Chem* 43:27-32.
114. Zeng, L. B., et al. 2011. Antioxidant activity and chemical constituents of essential oil and extracts of *Rhizoma Homalomenae*. *Food Chem* 125:456-463.
115. Zheng, W. and Wang, S.Y. 2001. Antioxidant activity and phenolic compounds in selected herbs. *J Agric Food Chem* 49: 5165-5170. 87.
116. Aruoma, O.I. 2003. Methodological consideration for characterization for potential antioxidant actions of bioactive components in plants foods. *Mutat Res.* 532: 9–20.
117. Mohammed, A.A. and Ibrahim, A.A. 2004. Pathological roles of reactive oxygen species and their defence mechanism. *Saudi Pharm J.* 12:1–18.
118. Bagchi, K. and Puri, S. 1998. Free radicals and antioxidants in health and disease. *East Mediterranean Health Jr.* 4:350–60.
119. Aruoma, O.I. 1994. Nutrition and health aspects of free radicals and antioxidants. *Food Chem Toxicol.* 32:671–83.
120. Cheeseman, K.H. and Slater, T.F. 1993. An introduction to free radicals chemistry. *Br Med Bull.* 49:481–93.
121. Rao, A.L., Bharani, M. and Pallavi, V. 2006. Role of antioxidants and free radicals in health and disease. *Adv Pharmacol Toxicol.* 7:29–38.
122. Esterbauer, H., Puhl, H. and Dieber-Rotheneder, M. 1991. Effect of antioxidants on oxidative modification of LDL. *Ann Med.* ; 23:573–81.
123. Neuzil, J., Thomas, S.R. and Stocker, R. 1997. Requirement for promotion, or inhibition of α -tocopherol of radical induced initiation of plasma lipoprotein lipid peroxidation. *Free Radic Biol Med.* 22:57–71.
124. Poppel, G.V. and Goldbohm, R.A. 1995. Epidemiologic evidence for β – carotene and cancer prevention. *Am J Clin Nutr*; 62:1393–5.
125. Glatthaar, B.E., Horing, D.H., M. Charan and Siroser, U. 1986. The role of ascorbic acid in carcinogenesis. *Adv Exp Med Biol.* 206: 357–77.
126. Sokol, R.J. 1988. Vitamin E deficiency and neurologic diseases. *Annu Rev Nutr.* 8: 351–73.
127. Ashok, B.T. and Ali, R. 1999. The aging paradox: Free radical theory of aging. *Exp Gerontol.* 34:293–303.
128. Sastre, J., Pellardo, F.V. and Vina, J. 1996. Glutathione, oxidative stress and aging. *Age.* 19:129–39.
129. Freeman, B.A. and Crapo, J.D. 1982. Biology of disease: Free radicals and tissue injury. *Lab Invest.* ; 47:412–26.
130. Hattori, Y., Nishigori, C., Tanaka, T., Ushida, K., Nikaido, O. and Osawa, T. 1997. 8 Hydroxy-2-deoxyguanosine is increased in epidermal cells of hairless mice after chronic ultraviolet B exposure. *J Invest Dermatol*; 89:10405–9.
131. Halliwell, B. 1995. How to characterize an antioxidant- An update. *Biochem Soc Symp.* 61:73–101.
132. Shi, H.L., Noguchi, N. and Niki, N. 1999. Comparative study on dynamics of antioxidative action of α -tocopheryl hydroquinone, ubiquinol and α -tocopherol, against lipid peroxidation. *Free Radic Biol Med.* 27: 334–46.
133. McHughen, A. 1990. "Flax (*Linum usitatissimum* L.): *In Vitro* Studies", *Biotechnology in Agriculture and Forestry*, Volume 10, pp 502-514.

134. Thompson, L.U., Chen, J.M., Li, T., Strasser-Weippl, K. and Goss, P.E. 2005. "Dietary flaxseed alters tumor biological markers in postmenopausal breast cancer". *Clin. Cancer Res.* 11 (10): 3828–35. doi:10.1158/1078-0432.CCR-04-2326.
135. Knight, J. 1998. Free radicals: Their history and current status in aging and disease. *Ann Clin Lab Sci.* 28:331–46.
136. Frie, B., Stocker, R. and Ames, B.N. 1988. Antioxidant defences and lipid peroxidation in human blood plasma. *Proc Natl Acad Sci.* 37: 569–71.
137. Rice-Evans, C.A. and Diplock, A.T. 1993. Current status of antioxidant therapy. *Free Radic Biol Med*; 15:77–96.
138. Krinsky, N.I. 1992. Mechanism of action of biological antioxidants. *Proc Soc Exp Biol Med.* 200: 248–54.
139. Sies, H. 1997. Oxidative stress: Oxidants and antioxidants. *Exp Physiol.* 82: 291–5.
140. Magnenat, J.L., Garganoam, M. and Cao, J. 1998. The nature of antioxidant defense mechanisms: A lesson from transgenic studies. *Environ Health Perspect*; 106:1219–28.
141. Banniste, J., Bannister, W. and Rotilio, G. 1987. Aspects of the structure, function, and applications of superoxide dismutase. *CRC Crit Rev Biochem.* 22: 111–80.
142. Johnson, F. and Giulivi, C. 2005. Superoxide dismutases and their impact upon human health. *Mol Aspects Med.* 26: 340–52.
143. Wuerges, J., Lee, J.W., Yim, Y.I., Yim, H.S., Kang, S.O. and DjinoVIC Carugo, K. 2004. Crystal structure of nickel containing superoxide dismutase reveals another type of active site. *Proc Natl Acad Sci.* 101: 8569–74.
144. Corpas, F.J., Barroso, J.B. and del Río, L.A. 2001. Peroxisomes as a source of reactive oxygen species and nitric oxide signal molecules in plant cells. *Trends Plant Sci.* 6: 145–50.
145. Chelikani, P., Fita, I. and Loewen, P.C. 2004. Diversity of structures and properties among catalases. *Cell Mol Life Sci.* 61: 192–208.
146. Gaetani, G., Ferraris, A., Rolfo, M., Mangerini, R., Arena, S. and Kirkman, H. 1996. Predominant role of catalase in the disposal of hydrogen peroxide within human erythrocytes. *Blood.* 87: 1595–9.
147. Meister, A. 1988. Glutathione metabolism and its selective modification. *J Biol Chem.* 263: 17205–8.
148. Fairlamb, A.H. and Cerami, A. 1992. Metabolism and functions of trypanothione in the Kinetoplastida. *Annu Rev Microbiol.* 46: 695–729.
149. Nassar, E., Mulligan, C., Taylor, L., Kerksick, C., Galbreath, M., Greenwood, M., et al. 2007. Effects of a single dose of N-Acetyl- 5-methoxytryptamine (Melatonin) and resistance exercise on the growth hormone/IGF-1 axis in young males and females. *J Int Soc Sports Nutr.* 4: 14.
150. Caniato, R., Filippini, R., Piovani, A., Puricelli, L., Borsarini, A. and Cappelletti, E. 2003. Melatonin in plants. *Adv Exp Med Biol.* 527: 593–7.
151. Reiter, R.J., Carneiro, R.C. and Oh, C.S. 1997. Melatonin in relation to cellular antioxidative defense mechanisms. *Horm Metab Res.* 29: 363–72.
152. Sanches Silva, A., Costa, D.D., Tabia-Albuquerque, G., Giovanna-Giuliana Buonocore, G.G., Ramos, F., Conceicao-Castilho, M., Machado, A.V. and Costa, H.S. 2014. Trends in the use of natural Antioxidants in active food Packaging: a review. *Food Additives and Contaminants: Part A* 31 (3):374-395.

153. Halliwell, B. 1997. Antioxidants and Human Disease A General Introduction. *Nutrition Reviews* 55 (1): S44-S49. DOI: 10.1111/j.1753-4887.1997.tb06100.x
154. Sen, C.K. 1995. Oxidants and antioxidant in exercise. *Journal of Applied Physiology* 79: 675-686.
155. Rebecca, S.Y., Wong, R.S.Y. and Radhakrishnan, A.K. 2012. Inhibition of In Vitro Human LDL Oxidation by Phenolic Antioxidants from Grapes and Wines. *Nutrition Review* 70 (9): 483-490. doi:10.1111/j.1753-4886. 2012.00512.x
156. Halliwell, B. 1994. Free radicals, antioxidant and human disease: curiosity, cause, or Consequence ? *The Lancet* 344 (8924):721-724.
157. Brewerv, M.S. 2011. Natural Antioxidants: sources, compounds, mechanisms of action and potential applications. *Comprehensive Reviews in Food Science and Food Safty* 10 (4): 221-247. DOI:10.1111/j.1541-4337.2011.00156.x
158. Shahidi, F. 2000. Antioxidants in food and Antioxidants LWT *Food Science and Technology* 4(3): 158-163.DOI:10.1002/1521-3803 (20000501)44:3<158::AID-FOOD 158>3.0.CO;2-L
159. Agarwal, A., Gupta, S. and Sikka, S. 2006. The role of free radicals and Antioxidants in reproduction. *Current Opinion in Obstetrics and Gynecology* 18 (3): 325-332. Doi: 10.1097/01.geo.0000193003.58158.4e
160. Priscilla, M. Clarkson, M.P. and Thompson, H.S. 2000. Antioxidants: what role do they play in physical activity and health ? *American Journal of Clinical Nutrition* 72 (2): 637s-646s.
161. Māheswari, R.K., Malhotra, J., Mohan, L. and Upadhyay, B. 2014. Exotic pepino:a shrub for Prophylactic Consequence and nutritional regime. *International Journal of Pharma Research and Health Sciences* 2 (1):42-48.
162. Rani, B., Singh, U. and Māheswari, R. 2013. Natural Antioxidants and their intrinsic worth for well-being. *Global Journal of Medical Research* XII (B-VII):54-69.

Corresponding author: Dr. Raaz K. Maheshwari, Department of Chemistry, SBRM Govt. P.G., College, Nagaur, Rajasthan, India
Email: drblj1682@gmail.com rkmgreenchem.jaipur@gmail.com